

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/14504

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☐

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/14504

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I, claim(s) 1-4, drawn to treatment or prevention of Alzheimer's disease by administering an inhibitor of phospholipase A2 activity.

Group II, claim(s) 5-7, drawn to use of a compound as in claim 5 to treat or prevent a physiological disorder associated with an excess of phospholipase A2.

The inventions listed as Groups I and II do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The treatment or prevention of Alzheimer's disease is a different condition from a physiological disorder associated with an excess of phospholipase A2 without a clear correlation (nexus) between the condition (excess phospholipase) and the disease. Thus, each method has its own separate special technical feature.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/41, 31/35, 31/355, 31/19, 31/165		A1	(11) International Publication Number: WO 95/17183
			(43) International Publication Date: 29 June 1995 (29.06.95)
(21) International Application Number: PCT/US94/14504 (22) International Filing Date: 14 December 1994 (14.12.94) (30) Priority Data: 08/173,544 23 December 1993 (23.12.93) US (71) Applicant: ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US). (72) Inventors: CLEMENS, James, Allen; 6350 Knyghton Road, Indianapolis, IN 46220 (US). SOFIA, Michael, Joseph; 3 Holly Lane, Lawrenceville, NJ 08648 (US). STEPENSON, Diane, Teresa; 9556 Mullet Court, Indianapolis, IN 46256 (US). (74) Agents: LAMMERT, Steven, R. et al.; Barnes & Thornburg, 1313 Merchants Bank Building, 11 South Meridian Street, Indianapolis, IN 46204 (US).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: USE OF PLA ₂ INHIBITORS AS TREATMENT FOR ALZHEIMER'S DISEASE			
(57) Abstract			
This invention provides methods for the treatment or prevention of Alzheimer's disease in a mammal which comprises administering to a mammal in need thereof an effective amount of an inhibitor of phospholipase A ₂ . This invention also provides a series of compounds which are useful as inhibitors of phospholipases A ₂ , especially cytosolic phospholipase A ₂ .			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LJ	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

USE OF PLA₂ INHIBITORS AS TREATMENT FOR
ALZHEIMER'S DISEASE

5 Alzheimer's disease is a degenerative disorder
of the human brain. Clinically, it appears as a
progressive dementia. Its histopathology is characterized
by degeneration of neurons, gliosis, and the abnormal
10 hallmarks include neurofibrillary tangles (paired helical
filaments) and amyloid deposits within the parenchyma and
cerebral vasculature.

While there is no general agreement as to the
chemical nature of neurofibrillary tangles, the major
15 constituent of both the amyloid plaque cores and the
amyloid of the congophilic angiopathy has been shown to be
a 4500 Dalton protein originally termed β -protein or
amyloid A4. Throughout this document this protein is
referred to as β -amyloid peptide or protein.

20 β -amyloid peptide is proteolytically derived
from a transmembrane protein, the amyloid precursor
protein. Different splice forms of the amyloid precursor
protein are encoded by a widely expressed gene. see, e.g.,
K. Beyreuther and B. Müller-Hill, Annual Reviews in
25 Biochemistry, 58:287-307 (1989). β -amyloid peptide
consists, in its longest forms, of 42 or 43 amino acid
residues. J. Kang, et al., Nature (London), 325:733-736
(1987). These peptides, however, vary as to their amino-
termini. C. Hilbich, et al., Journal of Molecular Biology,
30 218:149-163 (1991).

Because senile plaques are invariably surrounded
by dystrophic neurites, it was proposed early that β -
amyloid peptide is involved in the loss of neuronal cells
that occurs in Alzheimer's disease. B. Yankner and co-
35 workers were the first to demonstrate that synthetic β -
amyloid peptide could be neurotoxic in vitro and in vivo.

- 2 -

B.A. Yankner, et al., Science, 245:417 (1989); See also, N.W. Kowall, et al., Proceedings of the National Academy of Sciences, U.S.A., 88:7247 (1991). Other research groups, however, were unable to consistently demonstrate direct toxicity with β -amyloid peptide. See, e.g., Neurobiology of Aging, 13:535 (K. Kosik and P. Coleman, eds. 1992). Even groups receiving β -amyloid peptide from a common source demonstrate conflicting results. D. Price, et al., Neurobiology of Aging, 13:623-625 (1991) (and the references cited therein).

Because of the debilitating effects of Alzheimer's disease there continues to exist a need for effective treatments. This invention provides methods for the treatment of Alzheimer's disease in mammals.

Recent studies have begun to indicate that a major component of the pathology of Alzheimer's disease is chronic inflammation. See, J. Schnabel, Science, 260:1719-1720 (1993). Indeed, pathological investigations have demonstrated the presence of glial hyperactivity, acute phase proteins, and complement factors within affected areas of the brains of persons affected with Alzheimer's disease. Administration of nonsteroidal anti-inflammatory drugs appears to slow the advance of Alzheimer's disease. Id. Understanding this inflammatory component of Alzheimer's disease, therefore, will lead to advances in novel methods of treating patients suffering from this disease.

Inflammatory disorders account for a significant number of debilitating diseases. Inflammatory states, such as arthritis, psoriasis, asthma, and possibly atherosclerosis, stem from inflammatory reactions in the joints, skin, and blood vessels. It is generally believed that a central role in the inflammatory reaction is the production of phospholipid metabolites called eicosanoids. The eicosanoids represent a family of important mediators such as the leukotrienes, prostaglandins, lipoxins,

- 3 -

hydroxyeicosatetranoic acid, and thromboxanes. It is believed that the generation of eicosanoids is dependent on the availability of arachidonic acid which is liberated from phospholipids by the action of phospholipase A₂ (EC 3.1.1.4).

5 Phospholipase A₂ (PLA₂) is the common name for phosphatide 2-acylhydrolase, which catalyzes the hydrolysis of the sn-2-acyl ester bond of phosphoglycerides which results in the production of equimolar amounts of lysophospholipids and free fatty acids. see, E.A. Dennis, 10 "The Enzymes", Vol. 16, Academic Press, New York, (1983). Phospholipase A₂ enzymes are found in all living species and form a diverse family of enzymes. Over forty phospholipase A₂ enzymes have been structurally 15 characterized, and they show a high degree of sequence homology. J. Chang, et al., Biochemical Pharmacology, 36:2429-2436, (1987).

 The best characterized varieties of PLA₂ enzyme are the secreted forms, which are released into the 20 extracellular environment where they aid in the digestion of biological materials. The secreted forms have a molecular weight of about 12-15,000 (Chang, et al, supra). In contrast, cytosolic phospholipases A₂ are found in small amounts within the cell and play a key role in the 25 biosynthetic pathway leading to the formation of the platelet activating factors and the eicosanoids. D. Mobilio and L.A. Marshall, Annual Reports in Medicinal Chemistry, 24; 157-166, (1989).

 The cytosolic phospholipases A₂ have a molecular 30 weight of approximately 85,000 daltons. J.D. Clark, et al., Cell, 65:1043-1051 (1991). Free arachidonic acid is the rate limiting precursor for the production of eicosanoids and is liberated from its membrane phospholipid store by the action of cytosolic PLA₂. E.A. Dennis, Drug Development 35 and Research, 10:205-220, (1987). The same enzymatic step also produces lysophospholipids which may be converted to


- 4 -

platelet-activating factors. Thus, it is believed that cytosolic PLA₂ is central to the regulation of the biosynthetic pathways of potent lipid mediators of inflammation.

Due to the central role in the inflammatory component of Alzheimer's disease that appears to be played by cytosolic phospholipase A₂, it is desirable to identify and characterize new inhibitors of this enzyme.

The present invention describes a method for the treatment or prevention of Alzheimer's disease in a mammal which comprises administering to a mammal in need of said treatment an effective amount of an inhibitor of phospholipase A₂ activity or a pharmaceutically acceptable salt of said inhibitor.

The present invention also describes a method for the treatment or prevention of Alzheimer's disease in a mammal which comprises administering to a mammal in need of said treatment an effective amount of a compound of Formula I



I

wherein

R¹ is hydrogen, C₁-C₆ alkoxycarbonyl, C₁-C₆ alkoxy, C₂-C₆ alkanoyl, C₁-C₆ alkyl, or phenyl, said phenyl being optionally substituted with one or more halo substituents;

R² is hydroxy, C₁-C₆ alkoxy, hydrogen, or C₁-C₆ alkyl;

R³ is C₁-C₆ alkyl or hydrogen;

A is -O- or -CH₂-;

- 5 -



X^1 and X^2 are each $-CH_2-$ or taken together form $-CH=CH-$;

n is 0 to 6;

B is $-O-$, $-CH_2-$, or $-C(R^5R^6)-$;

5 where R^5 and R^6 are independently C_1-C_6 alkyl;

R^4 is phenyl, xanthenyl, tetrazolyl, or 3,4-dihydrobenzopyranyl, said phenyl, xanthenyl, or 3,4-dihydrobenzopyranyl being optionally substituted with one
10 or more substituents selected from the group consisting of C_1-C_6 alkyl, C_1-C_6 alkoxy, oxo, carboxy, C_1-C_6 alkoxy-carbonyl, carboxy- $(C_1-C_6$ alkoxy)-, carboxy- $(C_1-C_6$ alkyl)-, $NR^7R^8-C(O)-(C_1-C_6$ alkyl)-,



20 where R^7 and R^8 are independently hydrogen, C_1-C_6 alkyl, C_1-C_6 alkylsulfonyl, or phenylsulfonyl; or a pharmaceutically acceptable salt thereof.

In another embodiment this invention provides a method for the treatment or prevention of a condition associated with an excess of phospholipase A_2 activity
25 which comprises administering to a mammal in need thereof an effective amount of a compound of Formula I or a pharmaceutically acceptable salt thereof.

All temperatures stated herein are in degrees Celsius ($^{\circ}C$). All units of measurement employed herein are
30 in weight units except for liquids which are in volume units.

- 6 -

As used herein, the term "C₁-C₆ alkyl" represents, a straight or branched alkyl chain having from one to six carbon atoms. Typical C₁-C₆ alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, t-butyl, pentyl, neo-pentyl, hexyl and the like. The term "C₁-C₆ alkyl" includes within its definition the term "C₁-C₄ alkyl".

"Halo" represents chloro, fluoro, bromo or iodo. "C₁-C₆ alkoxy" represents a straight or branched alkyl chain having from one to six carbon atoms attached to an oxygen atom. Typical C₁-C₆ alkoxy groups include methoxy, ethoxy, propoxy, isopropoxy, butoxy, t-butoxy, pentoxy and the like. The term "C₁-C₆ alkoxy" includes within its definition the term "C₁-C₄ alkoxy".

"C₂-C₆ alkanoyl" represents a straight or branched alkyl chain having from one to five carbon atoms attached to a carbonyl moiety. Typical C₂-C₆ alkanoyl groups include ethanoyl, propanoyl, isopropanoyl, butanoyl, t-butanoyl, pentanoyl, hexanoyl, 3-methylpentanoyl and the like.


"C₁-C₄ alkoxycarbonyl" represents a straight or branched alkoxy chain having from one to four carbon atoms attached to a carbonyl moiety. Typical C₁-C₄ alkoxy-carbonyl groups include methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, t-butoxycarbonyl and the like.

The term "carboxy-(C₁-C₆ alkoxy)-" as used herein refers to a moiety of the structure



where m is 1-6, inclusive.

The term "carboxy-(C₁-C₆ alkyl)-" as used herein refers to a moiety of the structure



- 7 -

where m is 1 to 6, inclusive.

The term "hydroxy-protecting groups" as used herein refers to substituents of the hydroxy group commonly employed to block or protect the hydroxy functionality while reacting other functional groups on the compound. Examples of such hydroxy-protecting groups include methoxymethyl, benzyloxymethyl, methoxyethoxymethyl, 2-(trimethylsilyl)ethoxymethyl, methylthiomethyl, 2,2-dichloro-1,1-difluoroethyl, tetrahydropyranyl, phenacyl, cyclopropylmethyl, allyl, C₁-C₆ alkyl, 2,6-dimethylbenzyl, o-nitrobenzyl, 4-picolyl, dimethylsilyl, t-butyldimethylsilyl, levulinate, pivaloate, benzoate, dimethylsulfonate, dimethylphosphinyl, isobutyrate, adamantate and tetrahydropyranyl. Further examples of such groups may be found in T.W. Greene and P.G.M. Wuts, "Protective Groups in Organic Synthesis" (1991) at Chapter 3.

The term "leaving group" as used herein refers to a group of atoms that is displaced from a carbon atom by the attack of a nucleophile in a nucleophilic substitution reaction. The term "leaving group" as used in this document encompasses, but is not limited to, activating groups.

The term "activating group" as used herein refers a leaving group which, when taken with the carbonyl (-C=O) group to which it is attached, is more likely to take part in an acylation reaction than would be the case if the group were not present, as in the free acid. Such activating groups are well-known to those skilled in the art and preferably may be, for example, succinimidoxy, phthalimidoxy, benzotriazolyloxy or -O-CO-(C₄-C₇ alkyl).

The compounds used in the method of the present invention may have one or more asymmetric centers. As a consequence of these chiral centers, the compounds of the present invention occur as mixture of enantiomers.

- 8 -

racemates, racemic mixtures and as individual enantiomers. .
All asymmetric forms, individual isomers and combinations
thereof, are within the scope of the present invention.

As mentioned supra, the invention encompasses
5 methods employing the pharmaceutically acceptable salts of
the compounds defined by Formula I. A compound employed in
this invention can possess a sufficiently acidic, a
sufficiently basic, or both functional groups, and
accordingly react with any of a number of inorganic bases,
10 and inorganic and organic acids, to form a pharmaceutically
acceptable salt.

The term "pharmaceutically acceptable salt" as
used herein, refers to salts of the compounds of Formula I
which are substantially non-toxic to living organisms.
15 Typical pharmaceutically acceptable salts include those
salts prepared by reaction of the compounds of the present
invention with a pharmaceutically acceptable mineral or
organic acid or an inorganic base. Such salts are known as
acid addition and base addition salts.

20 Acids commonly employed to form acid addition
salts are inorganic acids such as hydrochloric acid,
hydrobromic acid, hydroiodic acid, sulfuric acid,
phosphoric acid, and the like, and organic acids such as
p-toluenesulfonic, methanesulfonic acid, oxalic acid,
25 p-bromophenylsulfonic acid, carbonic acid, succinic acid,
citric acid, benzoic acid, acetic acid, and the like.
Examples of such pharmaceutically acceptable salts are the
sulfate, pyrosulfate, bisulfate, sulfite, bisulfite,
phosphate, monohydrogenphosphate, dihydrogenphosphate,
30 metaphosphate, pyrophosphate, chloride, bromide, iodide,
acetate, propionate, decanoate, caprylate, acrylate,
formate, isobutyrate, caproate, heptanoate, propiolate,
oxalate, malonate, succinate, suberate, sebacate, fumarate,
maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate,
35 chlorobenzoate, methylbenzoate, dinitrobenzoate,
hydroxybenzoate, methoxybenzoate, phthalate, sulfonate,

- 9 -

xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, g-hydroxybutyrate, glycollate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate and the like. Preferred pharmaceutically acceptable acid addition salts are those formed with mineral acids such as hydrochloric acid and hydrobromic acid, and those formed with organic acids such as maleic acid and methanesulfonic acid.

Base addition salts include those derived from inorganic bases, such as ammonium or alkali or alkaline earth metal hydroxides, carbonates, bicarbonates, and the like. Such bases useful in preparing the salts of this invention thus include sodium hydroxide, potassium hydroxide, ammonium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate, potassium bicarbonate, calcium hydroxide, calcium carbonate, and the like. The potassium and sodium salt forms are particularly preferred.

It should be recognized that the particular counterion forming a part of any salt of this invention is usually not of a critical nature, so long as the salt as a whole is pharmacologically acceptable and as long as the counterion does not contribute undesired qualities to the salt as a whole.

The preferred methods of this invention employ those compounds in which:

R¹ is hydrogen, acetyl, propanoyl, methoxy, ethoxy, methoxycarbonyl, ethoxycarbonyl, methyl, ethyl, n-propyl, isopropyl, phenyl, monosubstituted phenyl, disubstituted phenyl, and trisubstituted phenyl;

R² is hydroxy, methoxy, ethoxy, hydrogen, methyl, ethyl, n-propyl, and isopropyl;

R³ is hydrogen, methyl, ethyl, n-propyl, isopropyl, and n-butyl;


n is 1 to 4;

- 10 -

B is -O-, -CH₂-, or -C(R⁵R⁶)-, where R⁵ and R⁶ are independently methyl, ethyl, or hydrogen; and

R⁴ is phenyl, xanthenyl, tetrazolyl, or 3,4-dihydrobenzopyranyl, said moieties being mono- or di-substituted with methyl, ethyl, propyl, isopropyl, butyl, methoxy, ethoxy, propoxy, butoxy, pentoxy, oxo, carboxy,

The compounds of this invention may be prepared according to standard methods known in the art. Many of the compounds employed in the methods of the present invention can be prepared by the methodology described in U.S. Patent 4,945,099, issued July 31, 1990, which is herein incorporated by reference. For example, the tetrazole compounds of Formula I may be prepared from the corresponding intermediate of Formula II



II

by any of a variety of standard methods. Generally, the nitrile is reacted with an azide reagent in a non-reactive solvent. Preferred conditions include the use of lithium or ammonium azide in dimethylformamide, sodium azide in diglyme and N,N-dimethylethanolamine hydrochloride, or tri-n-butyltin azide in a non-reactive solvent such as dimethoxyethane or tetrahydrofuran. Under the latter conditions, the reaction is generally heated at or near the reflux temperature of the reaction mixture. The transformation is generally complete under these conditions in 2-3 days. Other operable reaction conditions include the reaction of the nitrile of Formula II with an alkali metal azide such as sodium azide, ammonium chloride, and (optionally) lithium chloride in a non-reactive high-

- 11 -

boiling solvent such as N,N-dimethylformamide, preferably at temperatures from about 60°C. to about 125°C.

Alternatively, tri-n-butyltin azide or tetramethylguanidinium azide, in a solvent such as tetrahydrofuran, dimethoxyethane, diethoxyethane, or the like, may be used in place of the alkali metal azide, ammonium chloride, lithium chloride and N,N-dimethylformamide.


Similarly, the acids of this invention are prepared from the corresponding esters or nitriles. Hydrolysis of such esters or nitriles may be accomplished by any of a variety of acidic or basic conditions, preferably under aqueous conditions. Preferred methods involve the use of lithium hydroxide in a solvent mixture of acetone and water, sodium hydroxide in dioxane, or potassium hydroxide or potassium carbonate in a mixture of methanol and water. Under the former conditions, hydrolysis is generally complete in about 12-18 hours at temperatures from about 20-30°C whereas the latter reaction is usually complete in one hour at 20-30°C.

It is generally preferred, in compounds containing both a nitrile and an ester functionality, that the nitrile group be transformed into a tetrazole before hydrolysis of the ester.

Compounds of Formula I as well as intermediates in the preparation of the compounds of Formula I can be prepared by a number of synthetic routes as will be appreciated by skilled artisans depending upon the particular compound desired. For those compounds wherein A is -O-, the following scheme is generally applicable:

Scheme I

- 12 -



where L is a good leaving group such as halo, especially chloro, bromo or iodo, and R² is hydroxy or, preferably, a protected hydroxy group, such as benzyloxy.

The reaction of Scheme I is usually performed employing equimolar amounts of the two reactants although ratios other than equimolar amounts are completely operative. The reaction is best carried out in a nonreactive solvent such as a ketone, especially acetone or methyl ethyl ketone, or dimethylformamide, and in the presence of a base, preferably an alkali metal hydride or carbonate, preferably potassium carbonate. Especially when L is chloro, a catalyst such as potassium or sodium iodide may be added to increase the reaction rate. The reaction may be carried out at temperatures of about ambient temperature up to the boiling point of the reaction mixture, the former being preferred.

In the preferred case where the hydroxy group has been protected, the protecting group is removed following the coupling procedure described above. As will be appreciated by skilled artisans in the field, the means for deprotecting the hydroxy group will depend upon the choice of protecting group employed. In the preferred situation where a benzyl group is used, the benzyl group is removed by catalytic hydrogenation, for example, in the presence of 10% palladium on activated carbon in ethyl acetate, to provide the desired phenol. Usually this coupling reaction is performed before the final deprotecting of the R⁴ moiety; however, as will be appreciated, it is possible this sequence can be reversed depending on the functional groups involved. Thus, coupling as noted above


- 13 -

may, under certain circumstances well appreciated in the art, first involve transformation of the nitrile into 5-tetrazolyl followed by deprotection of the phenol.

A similar reaction protocol is found in Scheme

5 II:

Scheme II



10 where Q is bromo, chloro, iodo, mesyl, tosyl, or a similar leaving group. Aspects of this reaction scheme and all the variations thereof are generally the same as discussed above regarding Scheme I.

15 Other interconversions of compounds are readily apparent to skilled artisans. For example, when R⁴ is halo, compounds treated with cyanide, such as potassium or sodium cyanide, in a non-reactive solvent such as dimethylformamide, are transformed into cognates wherein R⁴ is -CN. The use of a catalytic amount of iodide is
 20 employed to speed the reaction. Such nitriles can then be converted into tetrazoles as described above, or hydrolyzed in the presence of a base, such as sodium or potassium hydroxide, in alcoholic water to provide the corresponding
 25 carboxylic acids. An alternate process for converting halides into nitriles involves the displacement by carbon anions in sodium amide and liquid ammonia as described in U.S. Patent 4,945,099.

30 Other transformations are also well known to those skilled in the art of organic chemistry. Carboxylic acids can be esterified by standard means, or converted to acid halides which are then reacted with amines to provide

- 14 -


the corresponding amides. Similarly, esters, amides, and nitriles may be hydrolyzed to the carboxylic acid. Nitriles can also be hydrolyzed to the primary amide by treatment with aqueous base.

5 The terms and abbreviations used in the instant examples have their normal meanings unless otherwise designated. For example "N" refers to normal or normality; "mmole" or "mmoles" refers to millimole or millimoles; "g" refers to gram or grams; "ml" means milliliter or
10 milliliters; "M" refers to molar or molarity; "eqv" refers to molar equivalents; "FDMS" refers to field desorption mass spectrometry; "MS" refers to mass spectrometry, and "NMR" refers to nuclear magnetic resonance.

15 The following examples further illustrate the preparation of the compounds of Formula I. These examples are illustrative only and are not intended to limit the scope of the invention in any way. In those compounds in which the terms "NMR" or "MS", or both, follow the
20 synthesis protocol, these terms indicate that the identity of the compounds was confirmed using nuclear magnetic resonance (NMR), mass spectrometry (MS) or both.

Example 1

25 N,N-Dimethyl-3-(2-(3-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)propoxy)phenyl)propionamide



- 15 -

5 A solution of 1.5 g of 3-(2-ethyl-4-bromo-5-benzyloxyphenoxy)propyl chloride and 0.5 g of tetrakis-(triphenylphosphine)palladium(0) in 70 ml of benzene was stirred with 15 ml of 2.0 M sodium carbonate. A solution of 1.1 g of 4-fluorophenyl boronic acid in 15 ml of ethanol was added. The mixture was heated at reflux for 16 hours. The mixture was cooled and diluted with ethyl acetate. The organic phase was washed with saturated ammonium chloride, washed with saturated sodium chloride, dried over sodium sulfate, and evaporated in vacuo. The residue was chromatographed on silica gel eluting with hexane/ethyl ether to provide 1.44 g (93%) of the desired title intermediate. NMR.

15 A mixture of 11 g of resorcinol, 8.8 g of 3-(2-ethyl-4-(4-fluorophenyl)-5-benzyloxyphenoxy)propyl chloride, and 13.8 g of potassium iodide in 150 ml of dimethylformamide was heated in an oil bath at 90°C for 24 hours. The mixture was cooled, diluted with water, and extracted with ethyl acetate. The organic phase was washed with water, washed with saturated sodium chloride, dried over sodium sulfate, and evaporated in vacuo. The residue was chromatographed on silica gel eluting with hexane/ethyl ether providing the title intermediate in 34% yield, NMR.

25 A solution of 375 mg of ethyl 3-(2-(3-(2-ethyl-4-(4-fluorophenyl)-5-benzyloxyphenoxy)propoxy)phenyl)-propionate in 25 ml of ethanol was mixed with 5 ml of 5.0 N sodium hydroxide and stirred 16 hours. The mixture was diluted with 1.0 N hydrochloric acid and extracted with 3:1 dichloromethane/isopropanol. The organic phase was washed with saturated sodium chloride, dried over sodium sulfate, and evaporated in vacuo providing the desired 3-(2-(3-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)propoxy)phenyl)propionic acid in 93% yield. NMR.


- 16 -

A solution of 3-(2-(3-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)propoxy)phenyl)propionic acid and several equivalents of thionyl chloride in dichloromethane was kept at room temperature for 3 hours, and then poured into a stirred solution of 40% dimethylamine in water. The organic layer was washed with aqueous hydrochloric acid, washed with saturated sodium chloride, dried over sodium sulfate, and evaporated in vacuo. The residue was chromatographed on silica gel eluting with ethyl acetate to provide the desired title product. NMR, MS.

15

Example 2

N-Methanesulfonyl-3-(2-(3-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)propoxy)phenyl)propionamide



20

A solution of 3-(2-(3-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)propoxy)phenyl)propionic acid and several equivalents of thionyl chloride in dichloromethane was maintained at room temperature for 3 hours to produce the acid chloride. To this acid chloride in tetrahydrofuran was added a suspension of 10 equivalents of N-lithiomethanesulfonamide in tetrahydrofuran at -5°C. The mixture was allowed to warm to room temperature,


25

- 17 -

diluted with aqueous hydrochloric acid, and extracted with ethyl acetate. The organic solution was dried and evaporated in vacuo. The residue was chromatographed on silica gel eluting with dichloromethane/methanol to provide the desired title intermediate in 37% yield. NMR.

Example 3


10 N-Phenylsulfonyl-3-(2-(3-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)propoxy)phenyl)propionamide




15 The title product was prepared by the procedure of Example 2 using N-lithiobenzenesulfonamide. The product was isolated by preparative C₁₈ reverse phase HPLC. NMR.

Example 4

20 Preparation of 2-phenyl-4-ethyl-5-[[6-(2H-tetrazol-5-yl)-6-methylheptyl]oxy]phenol



- 18 -



5 Synthesis of 1-Benzyl-2-phenyl-4-ethyl-5-(6-methyl-6-cyanoheptyloxy)benzene

A. Preparation of 4-benzyloxy-2-hydroxyacetophenone.

10

- 19 -

In a dry round-bottom flask under nitrogen, 2,4-dihydroxyacetophenone (15.2 g, 100 mmol) was dissolved in methyl ethyl ketone (400 ml) and dimethylsulfoxide (100 ml). To this solution were added benzyl bromide (17.0 g, 100 mmol) and potassium carbonate (27.6 g, 200 mmol). The reaction was heated to reflux and stirred for 15 hours. The methyl ethyl ketone was removed *in vacuo*, and the dimethylsulfoxide solution was diluted with ethyl acetate and washed several times with brine. The organic material was collected, dried (magnesium sulfate), filtered, and concentrated to provide a dark solid. The solid was recrystallized from hexane/toluene to provide the title benzyl ether as a tan solid (12.8 g, 55.7%); mp 143-144.5°C; NMR (CDCl₃) δ 12.77 (s, 1H), 7.70 (d, 1H, J = 7 Hz), 7.3-7.5 (m, 5H), 6.54 (d, 1H, J = 7 Hz), 6.53 (s, 1H), 5.11 (s, 2H), 2.58 (s, 3H).

Analysis for C₁₅H₁₂O₃:

Theory: C, 74.36; H, 5.82;

Found: C, 74.52; H, 5.97.

B. Preparation of 2-(6-methyl-6-cyanoheptyloxy)-4-benzyloxyacetophenone.

To a solution of 4-benzyloxy-2-hydroxyacetophenone (9.65 g, 42 mmol) in dimethylformamide (150 ml) were added the appropriate alkyl chloride (6.86 g, 40 mmol), potassium carbonate (10.6 g, 77 mmol), and potassium iodide (1.6 g, 9.6 mmol). The stirred reaction was heated to 90°C for 24 hours. The solids were removed by filtration, and the dimethylformamide was removed *in vacuo*. The residue was purified by Prep-500 HPLC, using a gradient of 5% ethyl acetate in hexane to 20% over 30 minutes as a mobile phase to yield the title ether as a clear oil (12.1 g, 79.8%); NMR (CDCl₃) δ 7.85 (d, 1H, J = 7.4 Hz), 7.3-7.5 (m, 5H), 6.60 (dd, 1H, J = 7.4, 1.8 Hz), 6.53 (d, 1H, J = 1.8 Hz),

- 20 -

5.12 (s, 2H), 4.04 (t, 2H, J = 5.3 Hz), 2.61 (s, 3H), 1.85-1.95 (m, 2H), 1.5-1.6 (m, 6H), 1.37 (s, 6H); IR (CHCl₃) 2943, 2238, 1601 cm⁻¹; MS (m/e) 379.

5 C. Preparation of 4-benzyloxy-2-(6-methyl-6-cyanoheptyloxy)ethylbenzene.

To a solution of 2-(6-methyl-6-cyanoheptyloxy)-4-benzyloxyacetophenone (12.1 g, 31.6 mmol) in carbon tetrachloride (30 ml) were added trifluoroacetic acid (44.4 g, 390 mmol) and triethylsilane (21.8 g, 188 mmol). The reaction was stirred at room temperature for 1.5 hours, then was worked-up by diluting with ethyl acetate and washing with aqueous sodium carbonate. The organic material was collected, dried (magnesium sulfate), filtered, and concentrated in vacuo. The residue was purified by Prep-500 HPLC using a 3% ethyl acetate in hexane to 5% grade over 15 minutes, then holding at 5%. Concentration of the appropriate fractions provided the desired title product (10.6 g, 91.5%) as a clear liquid. NMR (CDCl₃) δ 7.35-7.5 (m, 5H), 7.06 (d, 1H, J = 6.5 Hz), 6.53 (s, 1H), 6.52 (dd, 1H, J = 6.5, 2 Hz), 5.06 (s, 2H), 3.96 (t, 2H, J = 5.3 Hz), 2.60 (q, 2H, J = 6.3 Hz), 1.8-1.85 (m, 2H), 1.5-1.6 (m, 6H), 1.37 (s, 6H), 1.20 (t, 3H, J = 6.3 Hz).

25 D. Preparation of 1-bromo-2-benzyloxy-4-(6-methyl-6-cyanoheptyloxy)-5-ethylbenzene.

To a stirred solution of 4-benzyloxy-2-(6-methyl-6-cyanoheptyloxy)ethylbenzene (10.6 g, 28.9 mmol) in carbon tetrachloride (125 ml) was added N-bromosuccinimide (6.0 g, 33.3 mmol). Stirring was continued for 6 hours at room temperature. The mixture was then diluted with methylene chloride and washed with water. The organic material was collected, dried (magnesium sulfate), filtered, and concentrated in vacuo. The residue

- 21 -

was recrystallized from hexane/ethyl acetate to provide the title aryl bromide (12.6 g, 97.8%) as a pale yellow solid. NMR (CDCl₃) δ 7.35-7.5 (m, 5H), 7.22 (s, 1H), 6.50 (s, 1H), 5.17 (s, 2H), 3.90 (t, 2H, J = 5.3 Hz), 2.58 (q, 2H, J = 6.3 Hz), 1.75-1.85 (m, 2H), 1.50-1.65 (m, 6H), 1.37 (s, 6H), 1.18 (t, 3H, J = 6.3 Hz); IR (CHCl₃) 3020, 2981, 2946, 2238, 1662, 1600 cm⁻¹; MS (m/e) 444, 445, 446.

E. Representative procedures for the biaryl coupling reaction.

Method A

In a round-bottom flask, the appropriate aryl bromide (1 equivalent) was dissolved in benzene. To this solution were added Pd(PPh₃)₄ (10 mole %) and a 2.0 M aqueous solution of sodium carbonate (10 eq.). In a separate flask, the aryl boronic acid (2 eq.) was dissolved in ethanol. To the aryl boronic acid solution was added the the aryl bromide solution, and the mixture was heated to reflux and stirred for 16 hours. The mixture was diluted with ethyl acetate and washed with saturated aqueous ammonium chloride. The organic material was collected, dried (magnesium sulfate), filtered, and concentrated. The residue was purified by flash chromatography (6% ethyl acetate in hexane) to provide the desired biaryl.

Method B

A solution of the appropriate aryl bromide in tetrahydrofuran was cooled to -78°C. To this solution was added *tert*-butyl lithium (2 eq). The reaction was stirred at -78°C for 30 minutes, then a tetrahydrofuran solution of zinc chloride (1 eq) was added. The mixture was warmed to room temperature and stirred for 15 minutes. In a separate flask, a solution was prepared containing the appropriate aryl halide (1 eq) and Pd(PPh₃)₄ (10 mole%) in

- 22 -

tetrahydrofuran. This solution was added to the aryl zinc solution, and the mixture was stirred at room temperature for 2-18 hours. The reaction was diluted with ethyl acetate and washed with aqueous ammonium chloride. The organic material was dried (magnesium sulfate), filtered, and concentrated. The residue was purified by flash chromatography (6% ethyl acetate in hexane) to provide the desired biaryl.

10 F. Preparation of 1-benzyloxy-2-phenyl-4-ethyl-5-(6-methyl-6-cyanoheptyloxy)benzene.

This compound was prepared in 75% yield by Method A. NMR (CDCl₃) δ 7.60 (d, 2H, J = 6.5 Hz), 7.3-7.5 (m, 8H), 7.18 (s, 1H), 6.59 (s, 1H), 5.04 (s, 2H), 3.95 (t, 2H, J = 5.3 Hz), 2.63 (q, 2H, J = 6.3 Hz), 1.8-1.9 (m, 2H), 1.5-1.65 (m, 6H), 1.38 (s, 6H), 1.25 (t, 3H, J = 6.3 Hz); IR (CHCl₃) 3013, 2977, 2943, 2238, 1611, 1488 cm⁻¹; MS (m/e) 439.

20 Analysis for C₃₀H₃₅NO₂:

Calc: C, 81.59; H, 7.99; N, 3.17;

Found: C, 81.34; H, 8.18; N, 3.05.

25 To a solution of the nitrile (1 eq.) in diglyme were added N,N- dimethylethanolamine hydrochloride (2 eq.) and sodium azide (4 eq.). The suspension was heated to 130°C and stirred for up to 72 hours. The mixture was diluted with methylene chloride and acidified with dilute hydrochloride acid. The organic material was collected, dried (magnesium sulfate), filtered and concentrated in vacuo. The resulting material was dissolved in ethanol, and to this solution was added aqueous sodium hydroxide (4 eq.). This reaction was stirred at room temperature for 30 minutes, then the solvents were removed in vacuo. An HP-20 reverse phase MPLC system was used to purify the residue,

- 23 -

first using water as the mobile phase, then using 40% water, in methanol. The desired fractions were combined and concentrated *in vacuo*. The residue was then lyophilized to produce the tetrazole as its sodium salt.

5

2-Phenyl-4-ethyl-5-[6-(2H-tetrazol-5-yl)-6-methylheptyloxy]phenol sodium salt, 34.3% yield. NMR (DMSO-d₆) δ 7.55 (d, 2H, J = 6.5 Hz), 7.35 (t, 2H, J = 6.5 Hz), 7.20 (t, 1H, J = 6.5 Hz), 6.98 (s, 1H), 6.60 (s, 1H), 3.82 (t, 2H, J = 5.3 Hz), 2.65 (q, 2H, J = 6.3 Hz), 1.55-1.70 (m, 6H), 1.25-1.35 (m, 8H), 1.10 (t, 3H, J = 6.3 Hz); IR (KBr) 3192, 2970, 2937, 1617, 1488, 1453, 1214 cm⁻¹; MS (m/e) 439.

10

Analysis for C₂₃H₂₉N₄NaO₂·2H₂O:

15

Calc: C, 59.87; H, 7.16; N, 12.25;


Found: C, 60.28; H, 7.45; N, 12.07.

Example 5

20

2-[3-[3-[(5-Ethyl-2-hydroxy[1,1'-biphenyl]-4-yl)oxy]propoxy]-2-propylphenoxy]propanoic acid

- 24 -



A. Preparation of 2-propyl-1,3-dimethoxybenzene.

5

1,3-Dimethoxybenzene (20 g, 145 mmols) in 200 ml of dry tetrahydrofuran was cooled to -10°C . To this solution at -10°C was added n-butyllithium (100 ml of a 1.6 M solution in hexane, 160 mmols) over 20 minutes. The reaction was then stirred for 2.5 hours at 0°C . At 0°C , propyl iodide (24.65 g, 145 mmols) was added slowly over 15 minutes. When the addition was complete, the reaction was allowed to warm to room temperature and stirred overnight. After stirring overnight, the reaction was refluxed for 1.5 hours, then cooled to room temperature and quenched with ice. The tetrahydrofuran was removed under

- 25 -

vacuum, and the resulting aqueous layer was extracted several times with diethyl ether. The organic extract was dried over magnesium sulfate and filtered to give a clear oil after solvent removal (26.11 g). The oil was purified by vacuum distillation to provide the title intermediate (24.0 g, 92%).

Bp 80-82°C at 10 mm Hg.

NMR (CDCl₃) δ 7.16 (t, 1, J = 8.30 Hz), 6.58 (d, 2, J = 8.30 Hz), 3.85 (s, 6), 2.67 (t, 2, J = 7.57 Hz), 1.56 (m, 2), 0.99 (t, 3, J = 7.35 Hz).

B. Preparation of 2-propyl-1,3-dihydroxybenzene.

A mixture of solid 1,3-dimethoxy-2-propylbenzene (33.70 g, 190 mmoles) and solid pyridine hydrochloride (150 g, 1.30 mole) was warmed to 180°C. After 7.5 hours the reaction was cooled to 110°C and 50 ml of water was added slowly. After the reaction cooled to room temperature, it was diluted with 100 ml of water and extracted several times with ethyl acetate. The ethyl acetate extract was washed once with 2N hydrochloric acid and then dried over magnesium sulfate. Filtration and solvent removal gave 38.5 g of an orange solid. The title product was purified by recrystallization from dichloromethane providing 11.86 g (41%) of yellow crystals.

NMR (CDCl₃) δ 6.94 (t, 1, J = 8.10 Hz), 6.40 (d, 2, J = 8.10 Hz), 4.84 (s, 2), 2.63 (t, 2, J = 7.57 Hz), 1.62 (m, 2), 1.01 (t, 3, J = 7.33 Hz).

C. Preparation of ethyl 2-(2-propyl-3-hydroxyphenoxy)-propanoate.

- 26 -

Sodium hydride (1.08 g of a 60% oil dispersion, 27 mmoles) under an argon atmosphere was washed with 15 ml of dry hexane. The hexane supernatant was removed via syringe. Dry tetrahydrofuran (60 ml) was added to the sodium hydride and, with stirring at room temperature, the 2-propyl-1,3-dihydroxybenzene (4.08 g, 27 mmoles) was added as a 40 ml tetrahydrofuran solution. After stirring at room temperature for 25 minutes, the ethyl 2-bromopropionate (4.64 g, 26 mmoles) was added rapidly. After stirring at room temperature for 17 hours, the reaction was quenched with a saturated aqueous ammonium chloride solution and the tetrahydrofuran was removed under vacuum. The resulting aqueous mixture was extracted several times with ethyl acetate. The organic extract was dried over magnesium sulfate. Filtration and solvent removal gave an orange oil. This oil was purified by flash chromatography on silica gel eluting with 20% ethyl acetate/hexane. The desired title ester was obtained as a white solid (2.43 g, 36%).

TLC: R_f = 0.47 (30% ethyl acetate/hexane)
NMR ($CDCl_3$) δ 6.93 (dd, 1, J = 8.00 Hz), 6.45 (d, 1, J = 8.00 Hz), 6.30 (d, 1, J = 8.00 Hz), 5.77 (s, 1), 4.76 (q, 1, J = 6.76 Hz), 4.23 (q, 2, J = 7.02 Hz), 2.69 (m, 2), 1.63 (d, 3, J = 6.70 Hz), 1.60 (m, 2), 1.28 (t, 3, J = 7.50 Hz), 0.99 (t, 3, J = 7.50 Hz); IR (KBr) 3435, 2955, 2872, 1733, 1600, 1500, 1465 cm^{-1} ; Mass Spec. (FD) (m/z) 253 ($M^+ + 1$).

Analysis for $C_{14}H_{20}O_4$:
30 Calc: C, 66.65; H, 7.99;
 Found: C, 66.41; H, 8.04.

D. Preparation of ethyl 2-[3-[3-[(2-benzyloxy-1-bromo-5-ethyl-4-yl)oxy]propoxy]-2-propylphenoxy]propanoate.

- 27 -

Ethyl 2-(2-propyl-3-hydroxyphenoxy)propanoate was dissolved in methyl ethyl ketone (60 ml), and solid sodium iodide (20 g, 133 mmoles) was added. The reaction mixture was refluxed under an argon atmosphere for 18 hours. The reaction was cooled to room temperature, quenched with water, then extracted three times with diethyl ether. The organic extracts were combined, dried over magnesium sulfate, and filtered to give 6.27 g of a yellow oil.

A solution of ethyl 3,4-dihydro-7-hydroxy-8-propyl-2H-1-benzopyran-2-carboxylate (2.1 g, 8.1 mmoles) in dimethylformamide (5 ml) was added to a suspension of sodium hydride (324 mg, 8.1 moles, 60% oil dispersion) in 10 ml of dry dimethylformamide under a nitrogen atmosphere. After stirring the reaction mixture for 30 minutes, a mixture of the alkyl iodide (3.8 g, 8.1 mmoles) prepared above and 18-Crown-6 (110 mg, 0.4 mmole) was added. The reaction was stirred for 1.5 hours at room temperature. The reaction was quenched with water and then extracted several times with ethyl acetate. The organic material was dried over magnesium sulfate, filtered and concentrated under vacuum. The resulting product was purified by flash chromatography on silica gel eluting with 6% ethyl acetate/hexane to give the title intermediate as a clear oil (2.90 g, 68% yield).

TLC: R_f = 0.47 (30% ethyl acetate/hexane)

NMR (CDCl_3) δ 7.56-7.37 (m, 6), 7.12 (t, 1, J = 8.20 Hz), 6.62 (d, 1, J = 8.35 Hz), 6.59 (s, 1), 6.45 (d, 1, J = 8.31 Hz), 5.16 (s, 2), 4.80 (q, 1, J = 6.90 Hz), 4.26 (q, 2, J = 7.20 Hz), 4.18 (dd, 4, J = 5.91, 12.02 Hz), 2.80 (m, 2), 2.62 (q, 2, J = 7.47 Hz), 2.31 (m, 2), 1.69 (d, 3, J = 6.70 Hz), 1.65 (m, 2), 1.30 (t, 3, J = 7.20 Hz), 1.22 (t, 3, J = 7.54 Hz), 1.03 (t, 3, J = 7.35 Hz); IR (CHCl_3) 3015, 2967, 2930, 2780, 1752, 1595, 1500, 1464 cm^{-1} ; Mass Spec. (FAB) (m/z) 599 (M^+).

- 28 -

Analysis for $C_{32}H_{39}BrO_6$:

Calc: C, 64.11; H, 6.56; Br, 13.33;

Found: C, 64.01; H, 6.56; Br, 13.06.

- 5 E. Preparation of ethyl 2-[3-[3-[(2-benzyloxy-5-ethyl[1,1'-biphenyl]-4-yl)oxy]propoxy]-2-propylphenoxy]-propanoate.

10 Ethyl 2-[3-[3-[(2-benzyloxy-1-bromo-5-ethyl-4-yl)oxy]propoxy]-2-propylphenoxy]propanoate (1.3 g, 2.24 mmoles) was stirred in 40 ml of benzene under an argon atmosphere. To this solution was added

15 tetrakis(triphenylphosphine)palladium(0) (0.40 g, 0.35 mmole) and sodium bicarbonate (10 ml of a 2M aqueous solution). An ethanol solution (10 ml) of phenylboronic acid (1.3 g, 10.7 mmoles) was added to the above reaction mixture, and then the reaction mixture was refluxed for 21 hours. The reaction was cooled to room temperature,

20 quenched with a saturated aqueous ammonium chloride solution, diluted with water and then extracted with ethyl acetate. The organic layer was dried over magnesium sulfate and filtered. The filtrate was concentrated under vacuum providing 1.3 g of a brown solid. The solid was

25 dissolved in 20% ethyl acetate/hexane and filtered through 35 g of Merck 60 silica gel eluting with 500 ml of 20% ethyl acetate/hexane. The resulting 1.0 g of yellow oil was purified by flash chromatography on silica gel eluting with 18% ethyl acetate/hexane. The desired title ester was obtained in 47% yield as a clear oil.

30

TLC: R_f = 0.48 (30% ethyl acetate/hexane)

35 NMR ($CDCl_3$) δ 7.10 (d, 2, J = 8.06 Hz), 7.44 (m, 8), 7.27 (s, 1), 7.15 (t, 1, J = 8.14 Hz), 6.72 (s, 1), 6.66 (d, 1, J = 8.27 Hz), 6.48 (d, 1, J = 8.27 Hz), 5.11 (s, 2), 4.83 (q, 1, J = 6.71 Hz), 4.28 (m, 6), 2.78 (m, 4), 2.38 (m, 2), 1.72 (d, 3, J = 6.96 Hz), 1.69 (m, 2), 1.32 (t, 3, J = 7.29

- 29 -

Hz), 1.31 (t, 3, J = 7.30 Hz), 1.08 (t, 3, J = 7.36 Hz); IR (CHCl₃) 3015, 2966, 2930, 2880, 1750, 1594, 1488, 1464 cm⁻¹; Mass Spec. (FAB) (m/z) 597 (M⁺+1), 596 (M⁺).

Analysis for C₃₈H₄₄O₆:

5 Calc: C, 76.48; H, 7.43;
 Found: C, 76.42; H, 7.52.

10 F. Preparation of ethyl 2-[3-[3-[(5-ethyl-2-hydroxy[1,1'-biphenyl]-4-yl)oxy]propoxy]-2-propylphenoxy]-propanoate.

15 Hydrogen gas was bubbled for 15 minutes through a 10 ml ethyl acetate solution of ethyl 2-[3-[3-[(2-benzyloxy-5-ethyl[1,1'-biphenyl]-4-yl)oxy]propoxy]-2-propylphenoxy]propanoate containing 0.14 g of 10% palladium on activated carbon catalyst. A hydrogen atmosphere was maintained over the reaction mixture, and the reaction was stirred for 4 days. The reaction was filtered through a Celite® pad in a sintered glass funnel and the catalyst
20 was washed with ethyl acetate. The solvent was removed from the filtrate providing a clear oil. The oil was purified by flash chromatography on silica gel eluting with 20% ethyl acetate/hexane. The title intermediate was
25 obtained in 53% yield as a clear oil.

TLC: R_f = 0.36 (30% ethyl acetate/hexane)

30 NMR (CDCl₃) δ 7.43 (m, 5), 7.06 (d, 1, J = 8.84 Hz), 6.56 (s, 1), 6.37 (d, 1, J = 8.28 Hz), 5.20 (s, 1), 4.74 (q, 1, J = 6.73 Hz), 4.20 (m, 6), 2.71 (m, 2), 2.61 (q, 2, J = 7.58 Hz), 2.33 (t, 2, J = 6.05 Hz), 1.61 (d, 3, J = 6.94 Hz), 1.58 (m, 2), 1.25 (t, 3, J = 7.30 Hz), 1.19 (t, 3, J = 7.40 Hz), 0.96 (t, 3, J = 7.35 Hz); IR (CHCl₃) 3558, 3029, 3011, 2964, 2935, 2873, 1745, 1625, 1593, 1488, 1464 cm⁻¹; Mass Spec. (FAB) (m/z) 507 (M⁺+1), 506 (M⁺).

- 30 -

Analysis for $C_{31}H_{38}O_6$:

Calc:	C,	73.49;	H,	7.56;
Found:	C,	73.70;	H,	7.67.

- 5 G. Preparation of 2-[3-[3-[(5-ethyl-2-hydroxy[1,1'-biphenyl]-4-yl)oxy]propoxy]-2-propylphenoxy]propanoic acid.

10 A solution of ethyl 2-[3-[3-[(5-ethyl-2-hydroxy[1,1'-biphenyl]-4-yl)oxy]propoxy]-2-propylphenoxy]propanoate in 4 ml of dioxane was treated with 1.10 ml of 2N sodium hydroxide solution and stirred at room temperature. After 1.25 hours at room temperature, the dioxane was removed under vacuum and the remaining aqueous solution was diluted with water and acidified to pH 1 with 5N hydrochloric acid. The resulting suspension was extracted with ethyl acetate. The organic extract was dried over magnesium sulfate and filtered. The resulting white solid was recrystallized from toluene/hexane. The title product was crystallized from toluene/hexane and obtained as white tufts (0.582 g, 80%).

25 TLC: R_f = 0.21 (10% methanol/methylene chloride)
NMR ($CDCl_3$) δ 7.45 (m, 5), 7.09 (t, 1, J = 8.16 Hz), 7.03 (s, 1), 6.60 (d, 1, J = 8.28 Hz), 6.56 (s, 1), 6.42 (d, 1, J = 8.29 Hz), 4.79 (q, 1, J = 7.00 Hz), 4.20 (m, 4), 2.70 (m, 2), 2.62 (q, 2, J = 7.49 Hz), 2.33 (t, 2, J = 6.00 Hz), 1.67 (d, 3, J = 6.93 Hz), 1.56 (m, 2), 1.20 (t, 3, J = 7.39 Hz), 0.96 (t, 3, J = 7.30 Hz); IR (KBr) 3381, 2964, 2871, 1707, 1615, 1594, 1490, 1461 cm^{-1} ; Mass Spec. (FAB) (m/z) 479 ($M^+ + 1$), 478 (M^+).

Analysis for $C_{29}H_{34}O_6$:


Calc:	C,	72.78;	H,	7.16;
Found:	C,	73.39;	H,	7.29.

35

- 31 -

Example 6

8-Propyl-7-[3-[4-(4-fluorophenyl)-2-ethyl-5-hydroxyphenoxy]propoxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid



A. Preparation of ethyl 8-propyl-7-[3-[2-ethyl-4-(4-fluorophenyl)-5-benzyloxyphenoxy]propoxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylate.

Tetrakis(triphenylphosphine)palladium(0) (0.659 g, 0.6 mmole) and aqueous sodium carbonate solution (20 ml of a 2M solution) were added to a 30 ml benzene solution of ethyl 7-[3-[(2-benzyloxy-1-bromo-5-ethyl-4-yl)oxy]propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate (2.163 g, 3.5 mmole) under an argon atmosphere. The reaction was refluxed for 17 hours, then cooled to room temperature and extracted with ethyl acetate. The organic extract was dried over magnesium sulfate, filtered and the solvent removed under vacuum. The crude product was purified by Waters Prep 500 silica gel chromatography eluting with a gradient of 5% to 20% ethyl acetate/hexane over 50 minutes. The desired title biphenyl was obtained as a clear oil (1.722 g, 78%).

NMR (CDCl₃) δ 7.51 (m, 2), 7.32 (m, 5), 7.09 (m, 3), 6.83 (d, 1, J = 8.32 Hz), 6.62 (s, 1), 6.49 (d, 1, J = 8.50 Hz), 5.02 (s, 2), 4.75 (dd, 1, J = 4.10, 6.50 Hz), 4.22 (m, 6), 2.69 (m, 6), 2.25 (m, 4), 1.59 (m, 2), 1.30 (t, 3, J = 7.10)

- 32 -

Hz), 1.21 (t, 3, J = 7.42 Hz), 0.96 (t, 3, J = 7.33 Hz);
IR (CHCl₃) 3019, 2968, 1745, 1611, 1495 cm⁻¹; Mass Spec.
(FAB) (m/z) 627 (M⁺+1), 626 (M⁺), 536.

Analysis for C₃₉H₄₃O₆:

5 Calc: C, 74.74; H, 6.91; F, 3.03;
 Found: C, 74.98; H, 7.05; F, 3.39.

B. Preparation of ethyl 8-propyl-7-[3-[4-(4-
10 fluorophenyl)-2-ethyl-5-hydroxyphenoxy]propoxy]-3,4-
dihydro-2H-1-benzopyran-2-carboxylate.

Hydrogen gas was bubbled for 10 minutes through
a solution of ethyl 8-propyl-7-[3-[2-ethyl-4-(4-
15 fluorophenyl)-5-benzyloxy-phenoxy]propoxy]-3,4-dihydro-2H-
1-benzopyran-2-carboxylate (1.610 g, 2.57 mmols) in 30 ml
of ethyl acetate containing 1.0 g of 10% palladium on
activated carbon catalyst. The reaction was stirred at
room temperature under an atmosphere of hydrogen for 2
hours. The reaction mixture was filtered through a
20 Celite® pad in a sintered glass funnel and the catalyst was
washed with ethyl acetate. The solvent was removed from
the filtrate providing 1.242 g of a clear oil. The oil was
purified by flash chromatography on silica gel eluting with
20% ethyl acetate/hexane. The desired title phenol was
25 obtained in 74% yield (1.020 g) as a white solid.

TLC: R_f = 0.35 (30% ethyl acetate/hexane)

30 NMR (CDCl₃) δ 7.43 (m, 2), 7.16 (dd, 2, J = 5.97, 5.97 Hz),
6.98 (s, 1), 6.82 (d, 1, J = 8.44 Hz), 6.53 (s, 1), 6.46 (d,
1, J = 9.43 Hz), 5.07 (s, 1), 4.76 (m, 1), 4.21 (m, 6),
2.67 (m, 6), 2.26 (m, 4), 1.58 (m, 2), 1.29 (t, 3, J = 6.96
Hz), 1.91 (t, 3, J = 7.35 Hz), 0.96 (t, 3, J = 7.27 Hz);
IR (KBr) 3434, 2962, 2869, 1738, 1614, 1588, 1502 cm⁻¹;
Mass Spec (FAB) (m/z) 537 (M⁺+1), 536 (M⁺).

- 33 -

Analysis for C₃₂H₃₇O₆:

Calc:	C,	71.62;	H,	6.95;
Found:	C,	71.63;	H,	7.06.

5 C. Preparation of 8-propyl-7-[3-[4-(4-fluorophenyl)-2-ethyl-5-hydroxyphenoxy]propoxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid.

10 A dioxane (12 ml) solution of ethyl 8-propyl-7-[3-[4-(4-fluorophenyl)-2-ethyl-5-hydroxyphenoxy]propoxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylate (0.968 g, 1.8 mmoles) was treated with sodium hydroxide (2.71 ml of a 2N solution) and stirred at room temperature. After 2.5 hours at room temperature, the dioxane was removed from the
15 reaction mixture and the remaining material was diluted with water and acidified to pH 1.0 with 5N hydrochloric acid. The resulting white milky suspension was then stirred with ethyl acetate and subsequently extracted with ethyl acetate. The organic extract was dried over
20 magnesium sulfate, filtered and the solvent removed to give a white solid (1.098 g). The solid was recrystallized from ethyl acetate/hexane to give the title acid as white needle-like crystals (0.568 g, 62%).

25 TLC: R_f = 0.31 (10% methanol/methylene chloride)
NMR (CDCl₃) δ 7.42 (m, 2), 7.15 (dd, 2, J = 8.68), 6.98 (s, 1), 6.85 (d, 1, J = 8.30 Hz), 6.53 (s, 1), 6.52 (d, 1, J = 6.98 Hz), 4.77 (dd, 1, J = 3.63, 7.43 Hz), 4.18 (m, 4), 2.70 (m, 6), 2.27 (m, 4), 1.56 (m, 2), 1.19 (t, 3, J = 7.42
30 Hz), 0.95 (t, 3, J = 7.30 Hz); IR (KBr) 3421, 2959, 2871, 1706, 1615, 1500 cm⁻¹; Mass Spec (FAB) (m/z) 509 (M⁺+1), 508 (M⁺).

Analysis for C₃₀H₃₃O₆:

Calc:	C,	70.78;	H,	6.54;
Found:	C,	70.05;	H,	6.82.


35

- 34 -

Example 7

7-Carboxy-9-oxo-3-[3-(2-ethyl-5-hydroxy-4-phenylphenoxy)propoxy]-9H-xanthene-4-propanoic acid
disodium salt monohydrate

5



A mixture of 2-benzyloxy-1-phenyl-5-ethyl-4-(3-chloro-1-propyloxy)benzene (749 mg, 1.97 mmol), ethyl 7-carboethoxy-3-hydroxy-9-oxo-9H-xanthene-4-propanoate (729 mg, 1.97 mmol), potassium carbonate (1.36 g, 9.85 mmol) and potassium iodide (33 mg, 0.20 mmol) was refluxed for 24 hours. Dimethylsulfoxide (2 ml) was added and heating continued for 24 hours. The reaction mixture was cooled to room temperature, diluted with ethyl acetate, and washed once with water. The organic layer was dried over sodium sulfate, filtered and concentrated in vacuo to reveal a tan solid. This material was dissolved in ethyl acetate (30 ml) and the resulting solution purged with nitrogen. To this solution was added 10% palladium on carbon (120 mg) and the resulting suspension hydrogenated at 1 atmosphere of pressure. The solution was filtered and concentrated in vacuo to provide a colorless oil. This material was dissolved in a solution of 1:1 methanol/tetrahydrofuran (30 ml) and treated with 5N sodium hydroxide solution (2 ml) at room temperature for 18 hours. The resulting solution was extracted once with diethyl ether and the aqueous layer acidified with 5N hydrochloric acid solution. The resulting precipitate was collected via suction filtration.

- 35 -

This material was converted to the di-sodium salt and purified over HP-20 resin to provide 390 mg (56%) of the desired title product as a fluffy white solid: NMR (DMSO- d_6) 12.65 (s, 1H, -OH), 8.65 (s, 1H), 8.28 (dd, J = 8.5, 2.0 Hz, 1H), 8.01 (d, J = 8.9 Hz, 1H), 7.50 (m, 3H), 7.29 (t, J = 7.8 Hz, 2H), 7.17 (m, 2H), 6.93 (s, 1H), 6.89 (s, 1H), 4.26 (m, 4H), 3.12 (m, 2H), 2.47 (m, 2H), 2.23 (m, 2H), 1.10 (t, J = 7.4 Hz, 3H); MS-FAB m/e 627 (24, p), 605 (40), 583 (24), 331 (24), 309 (100); IR (KBr, cm^{-1}) 3419 (b), 2962, 1612, 1558, 1443, 1390, 1277, 1084.

Analysis for $C_{34}H_{28}O_9Na_2 \cdot H_2O$:


Calc:	C,	63.34;	H,	4.69;
Found:	C,	63.36;	H,	4.50.

15

Example 8

2-[2-Propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]phenoxy]benzoic acid

- 36 -



5 A. Preparation of 2-[2-propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-(phenylmethoxy)phenoxy]propoxy]phenoxy]-benzoic acid methyl ester.

10 A mixture of 2-benzyloxy-1-(4-fluorophenyl)-5-ethyl-4-(3-chloro-1-propyloxy)benzene (20.0 g, 50.2 mmols) and sodium iodide (75.3 g, 502 mmols) in 2-butanone (200 ml) was refluxed for 6 hours. The mixture was diluted with ether and washed once with water. The organic layer was dried

- 37 -

over sodium sulfate, filtered, and concentrated *in vacuo* to provide a colorless oil. This material was dissolved in dimethylformamide (100 ml) and treated with 2-(3-hydroxy-2-propylphenoxy)benzoic acid methyl ester (14.4 g, 50.2 mmoles) and potassium carbonate (20.8 g, 151 mmoles) at room temperature for 24 hours. This mixture was diluted with water and twice extracted with ether. The aqueous layer was separated and back-extracted once with ethyl acetate. The combined organic layers were dried over sodium sulfate, filtered, and concentrated *in vacuo* to provide a yellow oil. Silica gel chromatography provided 25.4 g (78%) of the desired title intermediate as a pale golden oil: NMR (CDCl₃) 7.91 (d, J = 7.8 Hz, 1H), 7.54 (d, J = 8.6 Hz, 1H), 7.52 (d, J = 8.5 Hz, 1H), 7.25-7.43 (m, 6H), 7.03-7.38 (m, 5H), 6.84 (d, J = 8.3 Hz, 1H), 6.71 (d, J = 8.1 Hz, 1H), 6.63 (s, 1H), 6.47 (d, J = 8.1 Hz, 1H), 5.03 (s, 2H), 4.24 (t, J = 5.7 Hz, 2H), 4.21 (t, J = 5.8 Hz, 2H), 3.86 (s, 3H), 2.69 (t, J = 7.8 Hz, 2H), 2.64 (t, J = 7.7 Hz, 2H), 2.34 (quintet, J = 6.0 Hz, 2H), 1.60 (hextet, J = 5.0 Hz, 2H), 1.22 (t, J = 7.5 Hz, 3H), 0.94 (t, J = 7.5 Hz, 3H); MS-FD m/e 648 (p); IR (CHCl₃, cm⁻¹) 2960, 1740, 1604, 1497, 1461, 1112.

Analysis for C₄₁H₄₁O₆F:

	Calc:	C,	75.91;	H,	6.37;
25	Found:	C,	76.15;	H,	6.45.

B. Preparation of 2-[2-propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]phenoxy]benzoic acid methyl ester.

2-[2-Propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-(phenylmethoxy)phenoxy]propoxy]phenoxy]benzoic acid methyl ester (33.0 g, 50.9 mmoles) was de-benzylated as described above for the preparation of Example 7 to provide 27.3 g (96%) of the title intermediate as an amber oil: NMR (CDCl₃) 7.90 (dd, J = 7.8, 1.7 Hz, 1H), 7.42 (m, 3H), 7.05-

- 38 -

7.23 (m, 4H), 6.99 (s, 1H), 6.84 (d, J = 8.1 Hz, 1H), 6.70 (d, J = 8.1 Hz, 1H), 6.55 (s, 1H), 6.46 (d, J = 8.1 Hz, 1H), 5.05 (s, 1H, -OH), 4.23 (m, 4H), 3.86 (s, 3H), 2.68 (t, J = 7.4 Hz, 2H), 2.62 (q, J = 7.5 Hz, 2H), 2.36 (quintet, J = 6.0 Hz, 2H), 1.60 (hextet, J = 7.7 Hz, 2H), 1.20 (t, J = 7.6 Hz, 3H), 0.94 (t, J = 7.4 Hz, 3H); MS-FD m/e 558 (p); IR (CHCl₃, cm⁻¹) 2965, 1727, 1603, 1496, 1458, 1306, 1112.

Analysis for C₃₄H₃₅O₆F:

10	Calc:	C,	73.10;	H,	6.31;
	Found:	C,	73.17;	H,	6.42.

C. Preparation of 2-[2-propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]phenoxy]benzoic acid sodium salt.

2-[2-Propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]phenoxy]benzoic acid methyl ester (21.5 g, 38.5 mmoles) was hydrolyzed as described above for the preparation of Example 7. The acid was converted to the sodium salt and purified as described in Example 7 to provide 16.7 g (77%) of the desired title product as a white amorphous solid: NMR (DMSO-d₆) 10.50 (bs, 1H, -OH), 7.51 (m, 3H), 7.20 (t, J = 7.4 Hz, 1H), 7.13 (m, 2H), 7.00 (m, 2H), 6.95 (s, 1H), 6.67 (dd, J = 8.2, 3.3 Hz, 2H), 6.62 (s, 1H), 6.26 (d, J = 8.2 Hz, 1H), 4.14 (t, J = 5.8 Hz, 2H), 4.02 (t, J = 5.7 Hz, 2H), 2.60 (t, J = 6.8 Hz, 2H), 2.47 (q, J = 7.3 Hz, 2H), 2.16 (t, J = 5.9 Hz, 2H), 1.45 (hextet, J = 7.5 Hz, 2H), 1.07 (t, J = 7.5 Hz, 3H), 0.81 (t, J = 7.4 Hz, 3H); MS-FAB m/e 568 (38, p + 1), 567 (100, p), 544 (86), 527 (77), 295 (65), 253 (45); IR (KBr, cm⁻¹) 3407 (b), 2962, 1603, 1502, 1446, 1395, 1239, 1112.


Analysis for C₃₃H₃₂O₆FNa:

35	Calc:	C,	69.95;	H,	5.69;	F,	3.35;
	Found:	C,	69.97;	H,	5.99;	F,	3.52.

- 39 -

Example 9


3-(2-(4-(2-Ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)-
 5 butyloxy)phenyl)propionic acid



A solution of 375 mg of ethyl 3-(2-(4-(2-ethyl-
 4-(4-fluorophenyl)-5-hydroxyphenoxy)-
 butyloxy)phenyl)propionate in 25 ml of ethanol was mixed
 with 5 ml of 5.0 N sodium hydroxide and stirred 16 hours.
 The mixture was diluted with 1.0 N hydrochloric acid and
 extracted with 3:1 dichloromethane/isopropanol. The
 organic phase was washed with saturated sodium chloride,
 dried over sodium sulfate, and evaporated in vacuo
 providing the desired title product in 72% yield. NMR.

Example 10

Preparation of 1-[5-ethyl-2-hydroxy-4-[[6-methyl-6-(1H-
 tetrazol-5-yl)heptyl]oxy]phenyl]ethanone



25


- 40 -

The title compound was prepared as described in U.S. Patent 4,945,099, issued July 31, 1990, which is herein incorporated by reference.

5

Example 11

Preparation of 2,4-diethyl-5-[[6-methyl-6-(1H-tetrazol-5-yl)heptyl]oxy]phenol



10


The title compound was prepared as described in U.S. Patent 4,945,099, issued July 31, 1990, which is herein incorporated by reference.

15

Example 12

Preparation of 3,4-dihydro-8-propyl-7-[3-(2-ethyl-5-hydroxy-4-ethoxy-phenoxy)propoxy]-2H-1-benzopyran-2-carboxylic acid

20



25

A. Preparation of 5-Ethyl-2,4-Dihydroxybenzaldehyde.

- 41 -

Dimethylformamide (250 ml) was cooled to 0°C under argon atmosphere. With stirring, phosphorous oxychloride (18.60 ml, 0.20 mole) was added slowly to the N,N-dimethylformamide. After several minutes at 0°C, the reaction was warmed to room temperature and methylene chloride (150 ml) was added to the reaction mixture to dissolve the solid. The reaction was subsequently recooled to 0°C. 4-Ethylresorcinol (25.0 g, 0.181 mole) was added to the reaction mixture as a 200 ml methylene chloride solution. After stirring at 0°C for 10 minutes., the reaction was warmed to room temperature and then refluxed for 16h. The reaction was cooled to room temperature, and a 100ml water solution of sodium acetate (50 g) was added slowly. This mixture was refluxed for 40 minutes then cooled to room temperature. The aqueous layer was washed several times with methylene chloride. The organics were combined and washed with 1N hydrochloric acid solution and brine the dried over magnesium sulfate. Filtration and solvent removal gave an orange solid which was recrystallized from toluene and hexane (17g, 56%)
TLC Rf = 0.39 (30% EtOAc/Hexane)

¹H NMR(CDCl₃) δ 11.30(s,1), 9.71(s,1), 7.29(s,1), 6.36(s(br),1), 2.61(q,2,J=7.48Hz), 1.25(t,3,J=7.54Hz)

B. Preparation of 1-(3-chloropropoxy-1-yl)-3-hydroxy-4-formyl-5-ethyl benzene.

A 190 ml dry tetrahydrofuran solution of 5-ethyl-2,4-dihydroxybenzaldehyde (8.00 g, 48.1 mmols), 3-chloropropanol (4.55 g, 48.1 mmols) and triphenylphosphine (12.62 g, 48.1 mmols) were stirred at room temperature. To this solution was added a 10 ml tetrahydrofuran solution

- 42 -

of diethyl azodicarboxylate (7.60 ml, 48.1 mmol). The reaction was stirred at room temperature for 17h after which the solvent was removed under vacuum. The crude material was adsorbed onto 125 g of 60 micron silica gel and then eluted through a plug 100 ml plug of silica gel with 1L of 30% ethyl acetate/hexane. The resulting yellow oil was then further purified by Waters Prep 500 chromatography on silica gel eluting with a solvent gradient of 5% to 30% ethyl acetate/hexane over 45 minutes. The desired product was obtained as a clear oil (7.03g, 61%).

TLC Rf = 0.47 (30% EtOAc/Hexane)

$^1\text{H NMR}$ (CDCl_3) δ 11.40 (s, 1), 9.71 (s, 1), 7.26 (s, 1), 6.42 (s, 1), 4.18 (t, 2, J=5.80 Hz), 3.77 (t, 2, J=6.28 Hz), 2.57 (q, 2, J=7.41 Hz), 2.30 (m, 2), 1.96 (t, 3, J=7.54 Hz)

IR (CHCl_3) 3021, 2971, 2937, 1643, 1586, 1494 cm^{-1}

Mass Spec (FD) m/e 243 (M^+)

Analysis for $\text{C}_{12}\text{H}_{15}\text{O}_3\text{Cl}$

Theory:	C, 59.33; H, 6.23; Cl, 14.61
Found	C, 59.24; H, 6.18; Cl, 14.69

C. Preparation of 1-Benzyloxy-3-[3-chloropropoxy]-4-ethyl-6-formyl benzene.

A suspension of hexane washed sodium hydride (2.40g of 60% oil dispersion, 60 mmol) in dry N,N-dimethylformamide was stirred under argon atmosphere at room temperature. A 50 ml dry N,N-dimethylformamide solution of 1-(3-chloropropoxy-1-yl)-3-hydroxy-4-formyl-5-ethyl benzene (6.92 g, 28.6 mmol) was added slowly to the NaH suspension, and this mixture was stirred for 30 minutes at room temperature. Benzyl bromide (9.78 g, 57.2 mmol) was added to the alkoxide solution and stirring was continued at room temperature. After three hours the reaction was

- 43 -

carefully quenched with saturated NH_4Cl solution and then ,
the reaction was diluted with water and extracted several
times with ethyl acetate. The organic extract was washed
with water and dried over magnesium sulfate. Filtration
5 and solvent removal gave a yellow solid. The solid was
purified by Waters Prep 500 chromatography using a silica
gel support and eluting with a solvent gradient of 5% to
40% ethyl acetate/hexane over a 45 minute period. The
desired product was obtained as a white solid (7.14 g,
10 75%).

TLC R_f = 0.34 (30% EtOAc/Hexane)

$^1\text{H NMR}(\text{CDCl}_3)$ δ 10.41(s,1), 7.68(s,1), 7.42(m,5), 6.49(s,1),
5.20(s,2), 4.17(t,2,J=5.78Hz), 3.78(t,2,J=6.22Hz),
15 2.57(q,2,J=7.53Hz), 2.29(m,2), 1.18(t,3,J=7.50Hz)

IR(CHCl_3) 3013, 2971, 2875, 1667, 1607, 1505, 1465 cm^{-1}

Mass Spec(FD) m/e 332(M^+)

20 Analysis for $\text{C}_{19}\text{H}_{21}\text{O}_3\text{Cl}$:

Theory C,68.57; H,6.36; Cl,10,65.

Found C,68.68; H,6.54; Cl,10,53. \approx

25 D. Preparation of 1-Benzyloxy-3-[3-chloropropoxy]-
4-ethyl-6-hydroxybenzene.

A solution of the aldehyde prepared supra in methylene
chloride (0.18 M) at room temperature was treated with m-
chloroperbenzoic acid (1.1 eqv), and the reaction was
30 stirred at room temperature. After 30 minutes. a
precipitate formed. The reaction was complete after 5
hours. The precipitate was removed by filtration. The
solvent was removed under vacuum and the resulting solid
35 was dissolved in tetrahydrofuran (0.28 M) and stirred
overnight with 2N sodium hydroxide (2.5 eqv).

- 44 -

Subsequently, the tetrahydrofuran was removed under vacuum, and the resulting aqueous mixture was diluted with water and acidified to pH 1 with 1N hydrochloric acid. The milky suspension was extracted several times with ethyl acetate. The organic extract was washed several times with saturated aqueous NaHCO₃ solution and then with brine. The organic layer was dried over magnesium sulfate. Filtration and solvent removal gave 8.70 g of a brown oil. The oil was purified by silica gel chromatography.

1-Benzyloxy-2-[3-chloropropoxy]-4-ethyl-6-hydroxybenzene was obtained in 66% yield from 1-Benzyloxy-2-[3-chloropropoxy]-4-ethyl-6-formylbenzene.

TLC Rf = 0.46 (30% EtOAc/Hexane)

¹HNMR(CDCl₃) δ 7.44(m,5), 6.79(s,1), 6.58(s,1), 5.25(s,1), 5.10(s,2), 4.03(t,2,J=5.77Hz), 3.78(t,3,J=6.36Hz), 2.56(q,2,J=7.56Hz), 2.23(m,2), 1.17(t,3,J=7.59Hz).

IR(CHCl₃) 3552, 3012, 2969, 2934, 1511, 1469 cm⁻¹

Mass Spec(FAB) m/e 320(M⁺)

Analysis for C₁₈H₂₁O₃Cl

Theory: C, 67.39; H, 6.60; Cl, 11.05.

Found: C, 67.09; H, 6.56; Cl, 10.82.

E. Preparation of 1-Benzyloxy-3-[3-chloropropoxy]-4-ethyl-6-ethoxy benzene.

A suspension of hexane washed sodium hydride (2.10 eqv) in dry N,N-dimethylformamide (1.3 M solution) was stirred under argon atmosphere at room temperature. A solution of the phenol in dry N,N-dimethylformamide (0.15 M) was added slowly to the sodium hydride suspension. The reaction was stirred at room temperature for 30 minutes. 18-Crown-6 was added to the reaction followed by the dropwise addition of

- 45 -

alkyl halide (5.0 eqv). After stirring at room temperature for several hours the reaction was quenched with saturated aqueous ammonium chloride solution, diluted with water and extracted with ethyl acetate. The ethyl acetate extract
5 was washed with water and then dried over magnesium sulfate. Filtration and solvent gave the crude product which was purified by silica gel flash chromatography.

1-Benzyloxy-2-[3-chloropropoxy]-4-ethyl-6-ethoxy benzene
10 was prepared in 77% yield as a white solid from 1-Benzyloxy-2-[3-chloropropoxy]-4-ethyl-6-hydroxy benzene and ethyliodide.

TLC Rf = 0.48 (30% EtOAc/Hexane)

¹HNMR(CDCl₃) δ 7.40(m,5), 6.80(s,1), 6.56(s,1), 5.16(s,2),
15 4.10(q,2,J=6.97Hz), 4.00(t,2,J=5.70Hz), 3.77(t, z,
J=6.73Hz), 2.60(q,2,J=7.50Hz), 2.21(m,2),
1.43(t,3,J=6.97Hz), 1.20(t,3,J=7.46Hz)

IR(CHCl₃) 3011, 2971, 2950, 2890, 1620, 1507, 1471 cm⁻¹
20

Mass Spec(FAB) m/e 348(M⁺)

Analysis for C₂₀H₂₅O₃Cl

Theory: C, 68.86; H, 7.22

Found: C, 69.35; H, 7.38
25

F. Preparation of Chromone

To a solution of 225 ml of absolute ethanol under argon
30 atmosphere and at room temperature was added 16.56 g of sodium metal over a 1 h. period. After all of the sodium metal was added the reaction mixture was refluxed for 1 hour, then cooled to room temperature. A mixture of 2,4-dihydroxyacetophenone (34.82g, 0.180 mole), diethyloxylate
35 (54.57ml, 0.41 mole), absolute ethanol (45 ml), and diethylether (45 ml) was added to the sodium ethoxide

- 46 -

5 solution over 25 minutes. The resulting deep maroon
reaction mixture was then refluxed for 2.5 hours and then
cooled to room temperature. The reaction mixture was
poured into approximately 600 ml of 1N hydrochloric acid
and then extracted several times with diethyl ether. The
ether was removed from the extract and the resulting gum
was dissolved in 135 ml of ethanol. To this solution was
then added 2.25 ml of concentrated hydrochloric acid and
subsequently refluxed for 45 minutes. The reaction was
10 cooled to room temperature and the ethanol was removed
under reduced pressure leaving a brown solid. This solid
was dissolved in ethyl acetate and washed once with water,
twice with saturated NaHCO₃, 1x with water and then dried
over magnesium sulfate. Filtration and solvent removal
15 gave 87g of a brown solid which was recrystallized from
ethyl acetate/petroleum ether. Recrystallization provided
24.07 g (48%) of a tan solid chromone.

TLC: Rf=0.27 (40% EtOAc/Hexane).

20 ¹H NMR (CDCl₃) δ 8.80 (s(br), 1), 7.98 (d, 1, J = 8.78Hz),
7.13 (d, 1, J = 8.78Hz), 7.13(s, 1), 4.47 (q, 2, J =
7.11Hz), 2.96 (t, 2, J = 7.25Hz), 1.73 (m, 2), 1.46 (t, 3,
J = 7.16Hz), 1.02 (t, 3, J = 7.11Hz).

25

G. Preparation of Ethyl 3,4-dihydro-8-propyl-2H-1-
benzopyran-2-carboxylate.

In a Parr bottle, chromone (12.07 g, 0.044 mole) was
30 dissolved in 210 ml of acetic acid. A catalyst (10%
palladium/activated carbon) (7.2 g) was added to this
solution and the bottle was pressurized with 52 psi of
hydrogen gas. The reaction was agitated for 23 hours. The
catalyst was removed by filtration through a Celite® pad
35 in a sintered glass funnel. The catalyst was washed with
ethyl acetate. The solvent was removed from the filtrate

- 47 -

and the resulting oil was azeotroped with toluene providing 12 g of brown oil. The material was purified on a Waters Prep 500 HPLC, equipped with silica gel cartridges, running a 5% to 40% ethyl acetate/hexane gradient over 50 minutes at a flow rate of 250 ml/min and collecting 500 ml fractions. The purified chroman was obtained as a pink oil (10 g, 86%).

TLC: Rf=0.50 (40% EtOAc/Hexane).

¹H NMR (CDCl₃) δ 6.73 (d, 1, J = 8.20Hz), 6.37 (d, 1, J = 8.20Hz), 4.78 (s(br), 1), 4.75 (m, 1), 4.25 (m, 2), 2.68 (m, 4), 2.16 (m, 2), 1.60 (m, 2), 1.29 (t, 3, J = 7.07Hz), 0.99 (t, 3, J = 7.34Hz).

H. Preparation of ethyl [3-[(1-benzyloxy-4-ethyl-2-ethoxy-5-yl)oxy]propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate.

To a solution of 1-benzyloxy-3-[3-chloropropoxy]-4-ethyl-6-ethoxy benzene (1.0 g, 2.87 mmol) in acetone (8.0 ml) and under argon atmosphere, added sodium iodide (4.31 g, 28.7 mmol). The reaction mixture was refluxed for 8h and then cooled to room temperature. The acetone was removed from the reaction mixture under vacuum, and the residue was dissolved in diethylether and washed with water. The ether extract was dried over magnesium sulfate and filtered. Solvent removal gave 1.09 g of the iodide as a yellow oil which solidified on standing at -4°C.

Under argon atmosphere and at room temperature, a mixture of ethyl 3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate (0.545 g, 2.06 mmol) and potassium carbonate (0.854 g, 6.18 mmol) in 4.0 ml of dry N,N-dimethylformamide was treated with a 4.0 ml N,N-dimethylformamide solution of the above prepared iodide.

- 48 -

After stirring at room temperature for 42 hours the reaction was quenched with water and then extracted several times with ethyl acetate. The ethyl acetate extract was washed with water and then dried over magnesium sulfate. Filtration and solvent removal gave 1.30 g of a yellow oil. The oil was purified by flash chromatography on silica gel eluting with 20% ethyl acetate/hexane. The desired coupled product (0.932 g) was obtained in 79% yield as a yellow oil.

TLC Rf = 0.48 (30% EtOAc/Hexane)

$^1\text{H NMR}(\text{CDCl}_3)$ δ 7.43(m,5), 6.85(d,1,J=8.37Hz), 6.82(s,1), 6.60(s,1), 6.50(d,1,J=8.37Hz), 5.13(s,2), 4.78(m,1), 4.26(m,2), 4.13(m,6), 2.67(m,6), 2.24(m,4), 1.62(m,2), 1.46(t,3,J=7.00Hz), 1.32(t,3, J=7.09Hz), 1.21(t,3,J=7.47Hz), 0.98(t,3,J=7.34Hz)

IR(CHCl₃) 3027, 3010, 2966, 2930, 2867, 1750, 1611, 1507, 1469 cm⁻¹

Mass Spec(FAB) m/e 486(M⁺)

Analysis for C₃₅H₄₄O₇

Theory:	C,72.89; H,7.69
Found:	C,72.85; H,7.40

I. Preparation of Ethyl [3-[(6-ethyl-4-ethoxy-3-hydroxy-1-yl) oxy]propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate.

To a solution of the benzyl ether in ethyl acetate or methanol (0.14 M solution) was added 10% palladium/activated carbon (15% w/w). Hydrogen gas was bubbled through this solution for 15 minutes. The reaction was then stirred at room temperature under an atmosphere of

- 49 -

hydrogen. After starting material was consumed, argon was bubbled through the reaction mixture for 15 minutes. The reaction mixture was filtered through a Celite® pad in a sintered glass funnel, and the catalyst was washed with ethyl acetate. The resulting crude product was purified by flash chromatography using silica gel as the solid support and eluting with an ethyl acetate/hexane mixture.

Ethyl [3-[(6-ethyl-4-ethoxy-3-hydroxy-1-yl) oxy]propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate was prepared from ethyl [3-[(1-benzyloxy-4-ethyl-2-ethoxy-5-yl)oxy]propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate in 87% yield as a white solid.

TLC Rf = 0.32 (30% EtOAc/Hexane)

¹HNMR(CDCl₃) δ 6.85(d,1,J=8.37Hz), 6.74(s,1), 6.62(s,1), 6.52(d,1,J=8.37Hz), 4.79(m,1), 4.19(m,8), 2.68(m,8), 2.26(m,4), 1.62(m,2), 1.45(t,3,J=6.96Hz), 1.33(t,3,J=7.14Hz), 1.21(t,3,J=7.54Hz), 1.00(t,3,J=7.33Hz)

IR(CHCl₃) 3540, 3026, 2965, 2934, 2873, 1750, 1611, 1509, 1492 cm⁻¹

Mass Spec(FD) m/e 486(M⁺)

Analysis for C₂₈H₃₈NO₇

Theory: C, 69.11; H, 7.87
Found: C, 69.00; H, 8.00

J. Preparation of 3,4-dihydro-8-propyl-7-[3-(2-ethyl-5-hydroxy-4-ethoxy-phenoxy)propoxy]-2H-1-benzopyran-2-carboxylic acid.

The ethyl ester was stirred in dioxane (0.14 M solution) at room temperature. This solution was treated with 3.0 eqv of sodium hydroxide (2N aq solution). The reaction was

- 50 -

stirred at room temperature for 2.5 hours and then the dioxane was removed under vacuum. The resulting residue was dissolved in water and acidified to pH 1 with 5N hydrochloric acid (a white ppt. forms). The aqueous mixture was extracted several times with ethyl acetate and then dried over magnesium sulfate. Filtration and solvent removal gave the crude product.

3,4-dihydro-8-propyl-7-[3-(2-ethyl-5-hydroxy-4-ethoxyphenoxy)propoxy]-2H-1-benzopyran-2-carboxylic acid was purified by recrystallization from ethyl acetate/hexane. The desired acid was obtained as white crystals (0.436 g, 71%)

TLC Rf = 0.49 (6.5/3.4/0.1 EtOAc/Hexane/AcOH)
 $^1\text{H NMR}$ (CDCl_3) δ 6.84(d,1,J=8.44Hz), 6.69(s,1), 6.56(s,1), 6.50(d,1,J=8.44Hz), 5.65(s(br),1), 4.77(dd,1,J=7.62, 3.72Hz), 4.10(m,6), 2.77(m,2), 2.62(m,4), 2.22(m,4), 1.54(m,2), 1.42(t,3,J=6.98Hz), 1.16(t,3,J=7.48Hz), 0.94(t,3,J=7.30Hz)

IR(KBr) 3215(br), 2956, 2930, 2870, 1706, 1613, 1589, 1516 cm^{-1}

Mass Spec(FD) m/e 458(M^+)
Analysis for $\text{C}_{26}\text{H}_{34}\text{O}_7$
Theory: C, 68.10; H, 7.47
Found: C, 68.13; H, 7.56


30

Example 13

Preparation of 7-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

35

- 51 -



A. Preparation of Chromone.

To a solution of 225 ml of ethanol (Anhydrous) under argon atmosphere. and at room temperature added 16.56 g of Na metal over a 1 hour. period. After all of the Na was added the reaction mixture was refluxed for 1 hour. then cooled to room temperature. A mixture of 2,4-dihydroxyacetophenone (34.82g, 0.180 mole), diethyloxylate (54.57ml, 0.41 mole), absolute ethanol (45 ml), and diethylether (45 ml) was added to the sodium ethoxide solution over 25 minutes. The resulting deep maroon reaction mixture was then refluxed for 2.5 hours and then cooled to room temperature. The reaction mixture was poured into approximately 600 ml of 1N hydrochloric acid and then extracted several times with diethyl ether. The ether was removed from the extract and the resulting gum was dissolved in 135 ml of ethanol. To this solution was then added 2.25 ml of concentrated hydrochloric acid and subsequently refluxed for 45 minutes. The reaction was cooled to room temperature and ethanol was removed under reduced pressure leaving a brown solid. This solid was dissolved in ethyl acetate and washed one time with water, two times with saturated sodium bicarbonate, one time with water and then dried over magnesium sulfate. Filtration and solvent removal gave 87g of a brown solid which was recrystallized from ethyl acetate/petroleum ether. Recrystallization provided 24.07 g (48%) of a tan solid chromone.

TLC: Rf=0.27 (40% EtOAc/Hexane).

- 52 -

¹H NMR (CDCl₃) δ 8.80 (s(br), 1), 7.98 (d, 1, J = 8.78Hz), 7.13 (d, 1, J = 8.78Hz), 7.13(s, 1), 4.47 (q, 2, J = 7.11Hz), 2.96 (t, 2, J = 7.25Hz), 1.73 (m, 2), 1.46 (t, 3, J = 7.16Hz), 1.02 (t, 3, J = 7.11Hz).

5

B. Preparation of Ethyl 3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate.

In a pressure bottle, the chromone (12.07 g, 0.044 mole) was dissolved in 210 ml of acetic acid. 10% palladium on activated carbon (7.2 g) catalyst was added to this solution and the bottle was pressurized with 52 psi of H₂ gas. The reaction was agitated for 23 hours. The catalyst was removed by filtration through a Celite® pad in a sintered glass funnel. The catalyst was washed with ethyl acetate. The solvent was removed from the filtrate and the resulting oil was azeotroped with toluene providing 12 g of brown oil. The material was purified on a Waters Prep 500 HPLC, equipped with silica gel cartridges, running a 5% to 40% ethyl acetate/hexane gradient over 50 minutes at a flow rate of 250 ml/minutes and collecting 500 ml fractions. The purified chroman was obtained as a pink oil (10 g, 86%).

TLC: R_f=0.50 (40% ethyl acetate/hexane).
¹H NMR (CDCl₃) δ 6.73 (d, 1, J = 8.20Hz), 6.37 (d, 1, J = 8.20Hz), 4.78 (s(br), 1), 4.75 (m, 1), 4.25 (m, 2), 2.68 (m, 4), 2.16 (m, 2), 1.60 (m, 2), 1.29 (t, 3, J = 7.07Hz), 0.99 (t, 3, J = 7.34Hz).

C. Ethyl 7-(3-chloropropoxy)-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate.

A solution (0.3M) of ethyl 3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate in dry N,N-dimethylformamide was stirred under argon atmosphere and at room temperature with solid potassium carbonate (1.75 eqv). To this

35

- 53 -

suspension added 1-bromo-2-chloropropane (2.5 eqv.). The reaction was stirred at room temperature for 20 hours and then quenched with water. The reaction mixture was extracted with ethyl acetate (three times), and the ethyl acetate extract was washed with water and then dried over magnesium sulfate. Filtration and solvent removal gave the crude product as an oil which was purified by flash chromatography on silica gel eluting with 15% ethyl acetate/hexane. Ethyl 7-(3-chloropropoxy)-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate was prepared in 72% yield.

^1H NMR (CDCl_3) δ 6.83(d,1,J=8.96Hz), 6.48 (d,1,J=8.96Hz), 4.77(t, 1, J=5.52Hz), 4.67(m,2), 4.10 (t,2,J=5.52Hz), 3.80 (t,2,J=5.50Hz) 2.70 (m,4), 2.26 (m,4), 1.6 (m,2), 1.28 (t,3,J=7.36Hz), 0.98 (t,3,J=6.44Hz)
IR (CHCl_3) 2963, 2933, 1749, 1728, 1612 cm^{-1}
Mass Spec (FAB) (m/z) 341($\text{M}^+\text{+H}$), 340(M^+)

D. Ethyl 7-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate.

To a solution of 2,4-dihydroxy-4-ethyl acetophenone in 5:1 methylethylketone and dimethylsulfoxide (0.45 M solution) at room temperature added the chloropropyl ether (1.0 eqv), potassium carbonate (1.75 eqv) and potassium iodide (0.20 eqv). The reaction was then refluxed for 20 hours. The reaction was then cooled to room temperature and quenched with water. The reaction mixture was extracted with ethyl acetate (three times) and this extract was washed with water and then dried over magnesium sulfate. Filtration and solvent removal gave a crude product which was purified by flash chromatography on silica gel eluting with 20% ethyl acetate/hexane. Ethyl 7-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate was prepared in 73% yield.

- 54 -

¹H NMR (CDCl₃) δ 7.43 (s,1), 6.81 (d,1,J=8.39Hz), 6.47 (d,1,J=8.39Hz), 6.42 (s,1), 4.75 (m,1), 4.24 (m, 4), 4.14 (t,2,J=5.98Hz), 2.64 (m,6), 2.58 (s,3), 2.35 (m,2), 2.20 (m,2), 1.55 (m,2), 1.29 (t,3,J=7.14Hz), 1.18 (t,3,J=7.47Hz), 0.93 (t,3,J=7.34Hz).

IR (CHCl₃) 2961, 2931, 2862, 1746, 1715, 1631, 1569 cm⁻¹

Mass Spec (FAB) (m/z) 485 (M⁺+H), 484 (M⁺)

Analysis for C₂₈H₃₆O₇:

Theory: C, 69.40; H, 7.49

Found: C, 70.23, H, 8.08

E. 7-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

A solution of ethyl 7-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate in dioxane (3.5 M solution) was treated with 2N sodium hydroxide (3.0 eqv) and stirred at room temperature. After stirring for 4 hours, the dioxane was removed from the reaction, and the remaining solution was diluted with water and acidified with 5N hydrochloric acid. The resulting milky solution was extracted with ethyl acetate. The ethyl acetate extract was dried over magnesium sulfate and filtered. Solvent removal gave a white solid. The solid was purified by flash chromatography on silica gel eluting with 50% ethyl acetate/hexane, and the resulting solid was crystallized from ethyl acetate and hexane. The desired acid was obtained in 47% yield.

¹H NMR (CDCl₃) δ 12.72 (s,1), 7.44 (s,1), 6.86 (d,1,J=8.30Hz), 6.51 (d,1,J=8.30Hz), 6.40 (s,1), 4.75 (dd,1,J=9.18Hz, 4.59Hz), 4.23 (t,2,J=5.74Hz), 4.15 (t,2,J=5.74Hz), 2.80 (m,1), 2.62 (m,2), 2.60 (s,3), 2.58 (m,2), 2.35 (m,2), 2.13 (m,1), 1.55 (m,2), 1.31 (t,3,J=6.90Hz), 1.20 (t,3,J=8.04Hz), 0.95 (t,3,J=8.04Hz)

- 55 -

IR (CCl₄) 3020, 3000, 2945, 3000, 1775, 1725, 1633, 1615
cm⁻¹

Mass Spec (FD) (m/z) 456 (M⁺)


Analysis for C₂₆H₃₂NO₇:

5 Theory: C, 68.40; H, 7.06
 Found: C, 68.61; H, 7.22

Example 14

10

Preparation of 2-[3-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)-propoxy]-2-propylphenoxy]butanoic acid.



15

A. Ethyl 2-[2-propyl-3-hydroxyphenoxy]butyrate.

 Sodium hydride (0.97 g of a 60% oil dispersion,
24 mmoles) under argon atmosphere was washed with 15 ml of
dry hexane. The hexane supernatant was removed via
20 syringe. Dry tetrahydrofuran (40 ml) was added to the
sodium hydride and with stirring at room temperature, the
dihydroxypropylbenzene (3.68 g, 24 mmoles) was added as a
40 ml tetrahydrofuran solution. After stirring at room
temperature for 25 minutes, the ethyl 2-bromo-propionate
25 (4.48 g, 23 mmoles) was added rapidly. After stirring at
room temperature for 17 hours, the reaction was quenched
with saturated aqueous ammonium chloride solution, and the
tetrahydrofuran was removed under vacuum. The resulting
aqueous mixture was extracted several times with ethyl
30 acetate. The organic extract was dried over magnesium
sulfate. Filtration and solvent removal gave an orange

- 56 -

oil. This oil was purified by Waters Prep 500 chromatography on silica gel eluting with 5 to 30% ethyl acetate/hexane gradient. The desired product was obtained as a clear oil (2.10 g, 33 %).

5 TLC Rf = 0.39 (30% EtOAc/Hexane, Silica gel)
 ^1H NMR (CDCl_3) δ 6.96 (t,1,J=8.2Hz), 6.45 (d,1,J=8.12Hz),
6.28 (d,1,J=8.12Hz), 4.88 (s,1), 4.59 (t,1,J=6.04Hz), 4.20
(q,2,J=7.52Hz), 2.69 (m,2), 2.02 (m,2), 1.63 (m,2), 1.24
(t,3,J=7.03Hz), 1.10 (t,3,J=7.43Hz), 0.99 (t,3,J=7.40Hz)
10 IR (CHCl_3) 3603, 3009, 2966, 2936, 2873, 1748, 1728, 1596
 cm^{-1}
Mass Spec (FAB) (m/z) 267 ($\text{M}^+\text{+H}$), 266 (M^+)

B. Ethyl 2-[2-propyl-3-(3-chloropropoxy)phenoxy]butyrate.

15 A solution (0.3M) of ethyl 2[2-propyl-3-hydroxyphenoxy]butyrate in dry N,N-dimethylformamide was stirred under argon atmosphere and at room temperature with solid potassium carbonate (1.75 eqv). To this suspension added 1-bromo-2-chloropropane (2.5 eqv.). The reaction was
20 stirred at room temperature for 20 hours and then quenched with water. The reaction mixture was extracted with ethyl acetate (three times), and the ethyl acetate extract was washed with water and then dried over magnesium sulfate. Filtration and solvent removal gave the crude product as an
25 oil which was purified by flash chromatography on silica gel eluting with 15% ethyl acetate/hexane. Ethyl 2-[2-propyl-3-(3-chloropropoxy)-phenoxy]butyrate was prepared in 85% yield.

^1H NMR (CDCl_3) δ 7.05 (t,1,J=8.26Hz), 6.55 (d,1,J=8.18Hz),
30 6.35 (d,1,J=8.27Hz), 4.60 (t,1,J=6.02Hz), 4.20
(q,2,J=7.13Hz), 4.11 (t,2,J=5.75Hz), 3.79
(t,2,J=6.36Hz), 2.72 (m,2), 2.26 (m,2), 2.01 (m,2), 1.59
(m,2), 1.25 (t,3,J=7.18Hz), 1.11 (t,3,J=7.39Hz),
0.97 (t,3,J=7.35Hz)
35 IR (CHCl_3) 3020, 2967, 2935, 2872, 1749, 1727, 1594 cm^{-1}
Mass Spec (FAB) (m/z) 343 ($\text{M}^+\text{+H}$), 342 (M^+)

- 57 -

Analysis for $C_{18}H_{27}O_4Cl$:

Theory: C, 63.06; H, 7.94; Cl, 10.34.

Found: C, 63.19; H, 7.84; Cl, 10.58.

- 5 C. Ethyl 2-[3-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)propoxy]-2-propylphenoxy]butanoate.

To a solution of 2,4-dihydroxy-4-ethyl acetophenone in 5:1 methylethylketone and dimethylsulfoxide (0.45 M soln) at room temperature was added ethyl 2-[2-propyl-3-(3-chloropropoxy)-phenoxy]butyrate (1.0 eqv), potassium carbonate (1.75 eqv) and potassium iodide (0.20 eqv). The reaction was then refluxed for 20 hours. The reaction was then cooled to room temperature and quenched with water. The reaction mixture was extracted with ethyl acetate (three times) and this extract was washed with water and then dried over magnesium sulfate. Filtration and solvent removal gave a crude product which was purified by flash chromatography on silica gel eluting with 20% ethyl acetate/hexane. Ethyl 2-[3-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)propoxy]-2-propylphenoxy]butanoate was prepared in 78% yield.

1H NMR ($CDCl_3$) δ 12.72 (s,1), 7.43 (s,1), 7.04 (t,1,J=8.29Hz), 6.55 (d,1,J=8.30Hz), 6.42 (s,1), 6.34 (d,1,J=8.30Hz), 4.58 (t,1,J=5.98Hz), 4.20 (m,6), 2.72 (m,2), 2.57 (s,3), 2.56 (m,2), 2.32 (m,2), 2.01 (m,2), 1.53 (m,2), 1.23 (t,3,J=7.06Hz), 1.18 (t,3,J=7.45Hz), 1.10 (t,3,J=7.38Hz), 0.94 (t,3,J=7.33Hz).

IR ($CHCl_3$) 2969, 2931, 1754, 1730, 1633, 1595 cm^{-1} Mass Spec (FD) (m/z) 486 (M^+)

- 30 Analysis for $C_{28}H_{38}O_7$:

Theory: C, 69.11; H, 7.87

Found: C, 69.08, H, 8.05

- 35 D. 2-[3-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)propoxy]-2-propylphenoxy]butanoic acid.

- 58 -

A solution of ethyl 2-[2-propyl-3-[[3-(4-acetyl-
2-ethyl-5-hydroxy-phenoxy)propyl]oxy]phenoxy]butanoate in
dioxane (3.5 M solution) was treated with 2N sodium
hydroxide (3.0 eqv) and stirred at room temperature. After
5 stirring for 4 hours, the dioxane was removed from the
reaction, and the remaining solution was diluted with water
and acidified with 5 N hydrochloric acid. The resulting
milky solution was extracted with ethyl acetate. The ethyl
acetate extract was dried over magnesium sulfate and
10 filtered. Solvent removal gave a white solid. The solid
was purified by crystallization from diethylether and
hexane. The desired acid was obtained in 69% yield.
¹H NMR (CDCl₃) δ 12.72 (s,1), 7.43 (s,1), 7.07
(t,1,J=8.28Hz), 6.58 (d,1,J=8.28Hz), 6.48 (s,1), 6.38
15 (d,1,J=8.28Hz), 4.63 (t,1,J=5.98Hz), 4.23 (t,2,J=6.00Hz),
4.17 (t,2,J=5.98Hz), 2.68 (m,2), 2.58 (s,3), 2.56 (m,2),
2.33 (m,2), 2.05 (m,2), 1.54 (m,2), 1.18 (t,3,J=7.42Hz),
1.12 (t,3,J=7.36Hz), 0.94 (t,3,J=7.29Hz)
IR (KBr) 2966, 2930, 2871, 1705, 1641, 1593 cm⁻¹
20 Mass Spec (FD) (m/z) 458 (M⁺)
Analysis for C₂₆H₃₄O₇:

Theory: C, 68.10; H, 7.47

Found: C, 68.01; H, 7.51

25

The biological activity of the compounds of
Formula I was evaluated employing an in vitro assay
measuring the ability of these compounds to inhibit the
activity of cytosolic phospholipase A₂. The assay was
30 performed essentially as described in R. Kramer, et al.,
Journal of Biological Chemistry, 266:5268-5272 (1991) with
the exception that varying amounts of the compound of
Formula I were added to the reaction mixture.

The substrate, sonicated liposomes containing
35 1-palmitoyl-2-[¹⁴C]arachidonoyl-sn-glycero-3-phosphocholine
([¹⁴C]PC, 55 mCi/mmol from NEN Research Products) and sn-

- 59 -

1,2-dioleoylglycerol (DG, Avanti Polar Lipids, Birmingham, Alabama) at a molar ratio of 2:1, was prepared as follows. [¹⁴C]PC (20 nmol, 1 x 10⁶ dpm, 50 µCi/ml in toluene/ethanol) and DG (10 nmol, 100 µg/ml in chloroform) were dried under nitrogen. The lipids were dispersed in 1 ml of 150 mM sodium chloride, 50 mM HEPES, pH 7.5 (assay buffer) by sonication at 4°C with a Microson™ probe-sonicator (Heat Systems Ultrasonics) for 4 x 15 seconds, with 45 second intervals. Bovine serum albumin (essentially fatty acid free, from a 100 mg/ml stock in water, Sigma) was added to a final concentration of 4 mg/ml. Samples to be assayed for cPLA₂ activity were incubated with 50 µl liposomes (0.5 nmol [¹⁴C]PC, 50,000 dpm containing 0.25 nmol of DG) and varying amounts of the compound of Formula I, in a total volume of 0.2 ml of assay buffer containing 1 mM calcium chloride and 1 mM 2-mercaptoethanol. Incubations were carried out at 37°C for 15 minutes and terminated by adding 2 ml of Dole's reagent (2-propanol/ heptane/0.5 M sulfuric acid, 40:10:1 containing 10 µg/ml of stearic acid).

After mixing, 1.2 ml of heptane and 1 ml of water were added. The mixtures were briefly vortexed and the upper phase transferred to tubes containing 2 ml of heptane and 150 mg of Bio-Sil (Bio-Rad Laboratories) activated at 130°C before use. The tubes were thoroughly vortexed and centrifuged (1000 x g for 5 minutes). The supernatants were decanted into scintillation vials. After addition of 10 ml of a liquid scintillation cocktail (Ready Protein®, Beckman) radioactivity was counted using a Beckman liquid scintillation counter Model LS 7000. High radioactive counts correlate with enzymatic activity.

Table I, *infra*, depicts the results of one such series of experiments. The first column provides the example number of the test inhibitor. The second column provides the concentration of the test compound (in

- 60 -

micromolar quantities) which inhibits fifty percent of the activity of cytosolic phospholipase A₂.

Table I

Inhibition of Cytosolic Phospholipase A₂

Example	Human Cytosolic
	PLA ₂ IC ₅₀ (μM)
2	18
3	17
4	39.4
5	12
6	9
8	12
9	35
10	43
11	54
12	28
13	13
14	14

Immunocytochemistry

10 Immunocytochemistry has demonstrated increased
numbers of reactive astrocytes containing cytosolic
phospholipase A₂ in the astrocytes of brains from patients
suffering from Alzheimer's disease. Immunocytochemistry was
performed on paraffin sections from human occipital cortex
15 of persons afflicted with Alzheimer's disease as well as
normal persons. In each case the tissue was fixed only
briefly (60-90 minutes) and then transferred to Tris-
buffered saline for several days prior to embedding. The
monoclonal antibody M12 was raised against purified cPLA₂

- 61 -

from U937 cells using standard techniques. Ascites were produced in BALB/c mice and antibodies were affinity-purified using Protein A Fast Flow™ resin. The antibody M12 recognizes the native form of cPLA₂ and is also a neutralizing antibody. A rabbit antiserum to glial fibrillary acidic protein (GFAP; Biogenex Labs, San Ramon, California) was used to label astrocytes.

Immunostaining of tissue sections (10 μm) utilized conventional immunoperoxidase techniques and employed the avidin-biotin peroxidase system (ABC, Vector Laboratories, Burlingame, California). For cPLA₂ localization, 0.1 mg/ml M12 antibody was used. Anti-GFAP was obtained as prediluted antisera. Dual localization was carried out by sequential immunostaining. An alkaline phosphatase-streptavidin system (Biogenex Labs) using Fast Red™ as chromagen was used to localize the rabbit antibody (GFAP) and nickel chloride-enhanced DAB (Vector Laboratories) was used to detect the peroxidase-labeled mouse anti-cPLA₂.

These immunochemistry studies demonstrated localization of cPLA₂ in protoplasmic astrocytes in the gray matter and provide further evidence for the importance of this cell type in inflammatory processes in the brain. Comparison of normal adult brains with those brains from persons afflicted with Alzheimer's disease evinces the role of cytosolic phospholipase A₂ in the inflammatory component of this disease.

Since the compounds employed in the present invention are effective inhibitors of cytosolic phospholipase A₂, these compounds are of value in the treatment of a wide variety of clinical conditions. This invention provides methods of treating or preventing Alzheimer's disease in a mammal which comprises administering to a mammal in need thereof an effective amount of a compound of Formula I.

- 62 -

The compounds of Formula I are usually administered in the form of pharmaceutical compositions. These compounds can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous, intravenous, intramuscular, and intranasal. These compounds are effective as both injectable and oral compositions. Such compositions are prepared in a manner well known in the pharmaceutical art and comprise at least one active compound.

In making the compositions employed in the present invention the active ingredient is usually mixed with an excipient, diluted by an excipient or enclosed within such a carrier which can be in the form of a capsule, sachet, paper or other container. When the excipient serves as a diluent, it can be a solid, semi-solid, or liquid material, which acts as a vehicle, carrier or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing for example up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions, and sterile packaged powders.

In preparing a formulation, it may be necessary to mill the active compound to provide the appropriate particle size prior to combining with the other ingredients. If the active compound is substantially insoluble, it ordinarily is milled to a particle size of less than 200 mesh. If the active compound is substantially water soluble, the particle size is normally adjusted by milling to provide a substantially uniform distribution in the formulation, e.g. about 40 mesh.

Some examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth,

- 63 -

gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl- and propylhydroxybenzoates; sweetening agents; and flavoring agents. The compositions of the invention can be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient by employing procedures known in the art.

The compositions are preferably formulated in a unit dosage form, each dosage containing from about 5 to about 100 mg, more usually about 10 to about 30 mg, of the active ingredient. The term "unit dosage form" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

The active compound is effective over a wide dosage range. For examples, dosages per day normally fall within the range of about 0.5 to about 30 mg/kg of body weight. In the treatment of adult humans, the range of about 1 to about 15 mg/kg/day, in single or divided dose, is especially preferred. However, it will be understood that the amount of the compound actually administered will be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response of the individual patient, and the severity of the patient's symptoms, and therefore the above dosage ranges are not intended to limit the scope of the invention in any way. In some instances dosage levels below the lower limit of the aforesaid range may be more than adequate, while in

- 64 -

other cases still larger doses may be employed without causing any harmful side effect, provided that such larger doses are first divided into several smaller doses for administration throughout the day.

5

Formulation Preparation 1

Hard gelatin capsules containing the following ingredients are prepared:

10	<u>Ingredient</u>	<u>Quantity</u> <u>(mg/capsule)</u>
	2-[2-Propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]phenoxy]benzoic acid	30.0
15	Starch	305.0
	Magnesium stearate	5.0

The above ingredients are mixed and filled into hard gelatin capsules in 340 mg quantities.

20

Formulation Preparation 2

A tablet formula is prepared using the ingredients below:

25	<u>Ingredient</u>	<u>Quantity</u> <u>(mg/tablet)</u>
	3-[5-methoxy-2-(4-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)-butyloxy)phenyl]propionic acid	25.0
30	Cellulose, microcrystalline	200.0
	Colloidal silicon dioxide	10.0
35	Stearic acid	5.0

- 65 -

The components are blended and compressed to form tablets, each weighing 240 mg.

Formulation Preparation 3

5

A dry powder inhaler formulation is prepared containing the following components:

	<u>Ingredient</u>	<u>Weight %</u>
10	8-Propyl-7-[3-[4-(4-fluorophenyl)-2-ethyl-5-hydroxyphenoxy]propoxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid	5
15	Lactose	95

The active mixture is mixed with the lactose and the mixture is added to a dry powder inhaling appliance.

Formulation Preparation 4

20

Tablets, each containing 30 mg of active ingredient, are prepared as follows:

	<u>Ingredient</u>	<u>Quantity (mg/tablet)</u>
25	2-phenyl-4-ethyl-5-[[6-(2H-tetrazol-5-yl)-6-methylheptyl]oxy]phenol	30.0 mg
	Starch	45.0 mg
30	Microcrystalline cellulose	35.0 mg
	Polyvinylpyrrolidone (as 10% solution in water)	4.0 mg
35	Sodium carboxymethyl starch	4.5 mg
	Magnesium stearate	0.5 mg

- 66 -

Talc	<u>1.0 mg</u>
Total	120 mg

5 The active ingredient, starch and cellulose are
passed through a No. 20 mesh U.S. sieve and mixed
thoroughly. The solution of polyvinylpyrrolidone is mixed
with the resultant powders, which are then passed through a
16 mesh U.S. sieve. The granules so produced are dried at
50-60°C and passed through a 16 mesh U.S. sieve. The
10 sodium carboxymethyl starch, magnesium stearate, and talc,
previously passed through a No. 30 mesh U.S. sieve, are
then added to the granules which, after mixing, are
compressed on a tablet machine to yield tablets each
weighing 150 mg.

15

Formulation Preparation 5

Capsules, each containing 40 mg of medicament
are made as follows:

20

<u>Ingredient</u>	<u>Quantity</u> <u>(mg/capsule)</u>
3,4-dihydro-8-propyl-7-(3-(2-ethyl-5- hydroxy-4-ethoxy-phenoxy)propoxy)-2H-1- 25 benzopyran-2-carboxylic acid	40.0 mg
Starch	109.0 mg
Magnesium stearate	<u>1.0 mg</u>
30 Total	150.0 mg

35 The active ingredient, cellulose, starch, and
magnesium stearate are blended, passed through a No. 20
mesh U.S. sieve, and filled into hard gelatin capsules in
150 mg quantities.

- 67 -

Formulation Preparation 6

Suppositories, each containing 225 mg of active
5 ingredient are made as follows:

	<u>Ingredient</u>	<u>Amount</u>
10	2,4-diethyl-5-[[6-methyl-6-(1H-tetrazol-5-yl)heptyl]oxy]phenol	25 mg
	Saturated fatty acid glycerides to	2,000 mg

The active ingredient is passed through a No. 60
mesh U.S. sieve and suspended in the saturated fatty acid
15 glycerides previously melted using the minimum heat
necessary. The mixture is then poured into a suppository
mold of nominal 2.0 g capacity and allowed to cool.

Formulation Preparation 7

20

Suspensions, each containing 50 mg of medicament
per 5.0 ml dose are made as follows:

	<u>Ingredient</u>	<u>Amount</u>
25	8-Propyl-7-[3-[4-(4-fluorophenyl)-2-ethyl-5-hydroxyphenoxy]propoxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid, ethyl ester	50.0 mg
30	Xanthan gum	4.0 mg
	Sodium carboxymethyl cellulose (11%)	
	Microcrystalline cellulose (89%)	50.0 mg
35	Sucrose	1.75 g
	Sodium benzoate	10.0 mg
	Flavor and Color	q.v.

- 68 -

Purified water to

5.0 ml

5 The medicament, sucrose and xanthan gum are blended, passed through a No. 10 mesh U.S. sieve, and then mixed with a previously made solution of the microcrystalline cellulose and sodium carboxymethyl cellulose in water. The sodium benzoate, flavor, and color are diluted with some of the water and added with stirring. Sufficient water is then added to produce the required

10 volume.

Formulation Preparation 8

15 Capsules, each containing 15 mg of medicament, are made as follows:

	<u>Ingredient</u>	<u>Quantity</u> <u>(mg/capsule)</u>
20	(E)-5-(3-carboxybenzoyl)-2-[[6-(4-methoxyphenyl)-5-hexenyl]oxy]-benzenepropanoic acid	15.0 mg
	Starch	407.0 mg
25	Magnesium stearate	<u>3.0 mg</u>
	Total	425.0 mg

30 The active ingredient, cellulose, starch, and magnesium stearate are blended, passed through a No. 20 mesh U.S. sieve, and filled into hard gelatin capsules in 560 mg quantities.

- 69 -

Formulation Preparation 9

An intravenous formulation may be prepared as follows:

5

	<u>Ingredient</u>	<u>Quantity</u>
10	7-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)-propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid	250.0 mg
	Isotonic saline	1000 ml

Formulation Example 10

15

A topical formulation may be prepared as follows:

	<u>Ingredient</u>	<u>Quantity</u>
20	3,4-dihydro-8-propyl-7-[3-(2-ethyl-5-hydroxy-4-ethoxy-phenoxy)propoxy]-2H-1-benzopyran-2-carboxylic acid	1-10 g
25	Emulsifying Wax	30 g
	Liquid Paraffin	20 g
	White Soft Paraffin	to 100 g

30 The white soft paraffin is heated until molten. The liquid paraffin and emulsifying wax are incorporated and stirred until dissolved. The active ingredient is added and stirring is continued until dispersed. The mixture is then cooled until solid.

35

- 70 -

Formulation Preparation 11

Sublingual or buccal tablets, each containing 10 mg of active ingredient, may be prepared as follows:

5		Quantity
	<u>Ingredient</u>	<u>Per Tablet</u>
	Active Ingredient	10.0 mg
10	Glycerol	210.5 mg
	Water	143.0 mg
	Sodium Citrate	4.5 mg
15	Polyvinyl Alcohol	26.5 mg
	Polyvinylpyrrolidone	<u>15.5 mg</u>
	Total	410.0 mg

20 The glycerol, water, sodium citrate, polyvinyl alcohol, and polyvinylpyrrolidone are admixed together by continuous stirring and maintaining the temperature at about 90°C. When the polymers have gone into solution, the solution is
25 cooled to about 50-55°C and the medicament is slowly admixed. The homogenous mixture is poured into forms made of an inert material to produce a drug-containing diffusion matrix having a thickness of about 2-4 mm. This diffusion matrix is then cut to form individual tablets having the
30 appropriate size.

Another preferred formulation employed in the methods of the present invention employs transdermal
35 delivery devices ("patches"). Such transdermal patches may be used to provide continuous or discontinuous infusion of

- 71 -

the compounds of the present invention in controlled amounts. The construction and use of transdermal patches for the delivery of pharmaceutical agents is well known in the art. See, e.g., U.S. Patent 5,023,252, issued June 11, 1991, herein incorporated by reference. Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.

Frequently, it will be desirable or necessary to introduce the pharmaceutical composition to the brain, either directly or indirectly. Direct techniques usually involve placement of a drug delivery catheter into the host's ventricular system to bypass the blood-brain barrier. One such implantable delivery system, used for the transport of biological factors to specific anatomical regions of the body, is described in U.S. Patent 5,011,472, issued April 30, 1991, which is herein incorporated by reference.

Indirect techniques, which are generally preferred, usually involve formulating the compositions to provide for drug latentiation by the conversion of hydrophilic drugs into lipid-soluble drugs or prodrugs. Latentiation is generally achieved through blocking of the hydroxy, carbonyl, sulfate, and primary amine groups present on the drug to render the drug more lipid soluble and amenable to transportation across the blood-brain barrier. Alternatively, the delivery of hydrophilic drugs may be enhanced by intra-arterial infusion of hypertonic solutions which can transiently open the blood-brain barrier.

30


- 72 -

Claims

5 1. A method for the treatment or prevention of
Alzheimer's Disease in a mammal which comprises
administering to a mammal in need of said treatment an
effective amount of an inhibitor of phospholipase A₂
10 activity or a pharmaceutically acceptable salt of said
inhibitor.

 2. A method as claimed Claim 1 wherein said
inhibitor of phospholipase A₂ activity is specific for
15 cytosolic phospholipase A₂.

 3. The use of a compound having the formula



20 wherein

R¹ is hydrogen, C₁-C₆ alkoxycarbonyl, C₁-C₆
alkoxy, C₂-C₆ alkanoyl, C₁-C₆ alkyl, or phenyl, said phenyl
being optionally substituted with one or more halo
substituents;

25 R² is hydroxy, C₁-C₆ alkoxy, hydrogen, or C₁-C₆
alkyl;

R³ is C₁-C₆ alkyl or hydrogen;

A is -O- or -CH₂-;

X¹ and X² are each -CH₂- or taken together form

30 -CH=CH-;


n is 0 to 6;

B is -O-, -CH₂-, or -C(R⁵R⁶)-;

- 73 -

where R⁵ and R⁶ are independently hydrogen or C₁-C₆ alkyl;

R⁴ is phenyl, xanthenyl, tetrazolyl, or 3,4-dihydrobenzopyranyl, said phenyl, xanthenyl, or 3,4-dihydrobenzopyranyl being optionally substituted with one or more substituents selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ alkoxy, oxo, carboxy, C₁-C₆ alkoxy-carbonyl, carboxy-(C₁-C₆ alkoxy)-, carboxy-(C₁-C₆ alkyl)-, NR⁷R⁸-C(O)-(C₁-C₆ alkyl)-,




where R⁷ and R⁸ are independently hydrogen, C₁-C₆ alkyl, C₁-C₆ alkylsulfonyl, or phenylsulfonyl; or a pharmaceutically acceptable salt or solvate thereof, in the preparation of a medicament useful for the treatment or prevention of Alzheimer's Disease.

4. The use of a compound as claimed in Claim 3 wherein said compound is selected from the group consisting of N,N-dimethyl-3-(2-(3-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)propoxy)phenyl)propionamide, N-methanesulfonyl-3-(2-(3-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)propoxy)phenyl)propionamide, N-phenylsulfonyl-3-(2-(3-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)propoxy)phenyl)propionamide, 8-propyl-7-(3-(4-(4-fluorophenyl)-2-ethyl-5-hydroxyphenoxy)propoxy)-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid, 2-[2-propyl-3-(3-(2-ethyl-4-(4-fluorophenyl)-5-

- 74 -

hydroxyphenoxy]propoxy]phenoxy]benzoic acid, 3-(2-(4-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)-butyloxy)phenyl)propionic acid, 2-[3-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)-propoxy]-2-propylphenoxy]butanoic acid, 5 7-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid, 2-phenyl-4-ethyl-5-[[6-(2H-tetrazol-5-yl)-6-methylheptyl]oxy]phenol, 1-[5-ethyl-2-hydroxy-4-[[6-methyl-6-(1H-tetrazol-5-yl)heptyl]oxy]phenyl]ethanone, 2,4- 10 diethyl-5-[[6-methyl-6-(1H-tetrazol-5-yl)heptyl]oxy]phenol, 2-[3-[3-[(5-ethyl-2-hydroxy[1,1'-biphenyl]-4-yl)oxy]propoxy]-2-propylphenoxy]propanoic acid, 7-carboxy-9-oxo-3-[3-(2-ethyl-5-hydroxy-4-phenylphenoxy)propoxy]-9H-xanthene-4-propanoic acid, 3,4-dihydro-8-propyl-7-[3-(2- 15 ethyl-5-hydroxy-4-ethoxy-phenoxy)propoxy]-2H-1-benzopyran-2-carboxylic acid, 2-[3-[3-[(5-ethyl-2-hydroxy[1,1'-biphenyl]-4-yl)oxy]propoxy]-2-propylphenoxy]propanoic acid, 7-carboxy-9-oxo-3-[3-(2-ethyl-5-hydroxy-4-phenylphenoxy)propoxy]-9H-xanthene-4-propanoic acid, and 20 3,4-dihydro-8-propyl-7-[3-(2-ethyl-5-hydroxy-4-ethoxy-phenoxy)propoxy]-2H-1-benzopyran-2-carboxylic acid, or a pharmaceutically acceptable salt or solvate of any of these compounds.

25 5. The use of a compound having the formula



wherein

30 R¹ is hydrogen, C₁-C₆ alkoxy carbonyl, C₁-C₆ alkoxy, C₂-C₆ alkanoyl, C₁-C₆ alkyl, or phenyl, said phenyl

- 75 -

being optionally substituted with one or more halo substituents;

R^2 is hydroxy, C_1 - C_6 alkoxy, hydrogen, or C_1 - C_6 alkyl;

R^3 is C_1 - C_6 alkyl or hydrogen;

A is -O- or -CH₂-;


X^1 and X^2 are each -CH₂- or taken together form -CH=CH-;

n is 0 to 6;

B is -O-, -CH₂-, or -C(R^5 R^6)-;

where R^5 and R^6 are independently hydrogen or C_1 - C_6 alkyl;

R^4 is phenyl, xanthenyl, tetrazolyl, or 3,4-dihydrobenzopyranyl, said phenyl, xanthenyl, or 3,4-dihydrobenzopyranyl being optionally substituted with one or more substituents selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 alkoxy, oxo, carboxy, C_1 - C_6 alkoxycarbonyl, carboxy-(C_1 - C_6 alkoxy)-, carboxy-(C_1 - C_6 alkyl)-, NR^7R^8 -C(O)-(C₁-C₆ alkyl)-,



where R^7 and R^8 are independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkylsulfonyl, or phenylsulfonyl; or a pharmaceutically acceptable salt or solvate thereof, in the preparation of a medicament useful for the treatment or prevention of a physiological disorder associated with an excess of phospholipase A₂.

- 76 -

6. The use of a compound as claimed in Claim 5 wherein said compound is selected from the group consisting of N,N-dimethyl-3-(2-(3-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)propoxy)phenyl)propionamide, N-methanesulfonyl-3-(2-(3-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)propoxy)phenyl)propionamide, N-phenylsulfonyl-3-(2-(3-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)propoxy)phenyl)propionamide, 8-propyl-7-[3-[4-(4-fluorophenyl)-2-ethyl-5-hydroxyphenoxy]propoxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid, 2-[2-propyl-3-[3-[2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy]propoxy]phenoxy]benzoic acid, 3-(2-(4-(2-ethyl-4-(4-fluorophenyl)-5-hydroxyphenoxy)-butyloxy)phenyl)propionic acid, 2-[3-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)-propoxy]-2-propylphenoxy]butanoic acid, 7-[3-(4-acetyl-2-ethyl-5-hydroxyphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid, 2-phenyl-4-ethyl-5-[[6-(2H-tetrazol-5-yl)-6-methylheptyl]oxy]phenol, 1-[5-ethyl-2-hydroxy-4-[[6-methyl-6-(1H-tetrazol-5-yl)heptyl]oxy]phenyl]ethanone, 2,4-diethyl-5-[[6-methyl-6-(1H-tetrazol-5-yl)heptyl]oxy]phenol, 2-[3-[3-[(5-ethyl-2-hydroxy[1,1'-biphenyl]-4-yl)oxy]propoxy]-2-propylphenoxy]propanoic acid, 7-carboxy-9-oxo-3-[3-(2-ethyl-5-hydroxy-4-phenylphenoxy)propoxy]-9H-xanthene-4-propanoic acid, 3,4-dihydro-8-propyl-7-[3-(2-ethyl-5-hydroxy-4-ethoxy-phenoxy)propoxy]-2H-1-benzopyran-2-carboxylic acid, 2-[3-[3-[(5-ethyl-2-hydroxy[1,1'-biphenyl]-4-yl)oxy]propoxy]-2-propylphenoxy]propanoic acid, 7-carboxy-9-oxo-3-[3-(2-ethyl-5-hydroxy-4-phenylphenoxy)propoxy]-9H-xanthene-4-propanoic acid, and 3,4-dihydro-8-propyl-7-[3-(2-ethyl-5-hydroxy-4-ethoxy-phenoxy)propoxy]-2H-1-benzopyran-2-carboxylic acid, or a pharmaceutically acceptable salt or solvate of any of these compounds.

35

- 77 -

7. The use of a compound as claimed in Claim 5 , wherein the physiological disorder associated with an excess of of phospholipase A₂ is associated with an excess of cytosolic phospholipase A₂.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/14504

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 31/41, 31/35, 31/355, 31/19, 31/165
US CL : 514/381, 454, 455, 456, 457, 458, 568, 570, 571, 622

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : none

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
none

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
APS and CAS: compounds of the claims with Alzheimer's, phospholipase A2, PLA2

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A, 4,814,339 (ROTONDO) 21 MARCH 1989, column 1, lines 5-61.	1-4
Y	US, A, 4,788,304 (MARSHALL ET AL.) 29 NOVEMBER 1988, column 1, line 4 - column 3, line 15.	5-7
Y	US, A, 5,070,207 (SCHIEHSER ET AL.) 3 DECEMBER 1991, column 1, line 4 - column 6, line 28.	5-7
Y,P	US, A, 5,290,817 (PETRAITIS) 01 MARCH 1994 column 1, line 6 - column 5, line 16.	5-7

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* "A"	document defining the general state of the art which is not considered to be of particular relevance	* "T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* "E"	earlier document published on or after the international filing date	* "X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* "L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* "Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* "O"	document referring to an oral disclosure, use, exhibition or other means	* "A"	document member of the same patent family
* "P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

07 APRIL 1995

Date of mailing of the international search report

21 APR 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized Officer
KIMBERLY R. JORDAN

Telephone No. (703) 305-1235